

AROUND THE GROTHENDIECK ANABELIAN SECTION CONJECTURE

MOHAMED SAÏDI

ABSTRACT. This paper is around the topics I discussed in the lecture I gave at the Isaac Newton Institute in Cambridge, July 2009, in the Introductory Workshop. This paper can be read as a companion to my paper [Saïdi], where detailed proofs can be found.

CONTENTS

- §0. Introduction
- §1. Generalities on Arithmetic Fundamental Groups and Sections
- §2. Grothendieck Anabelian Section Conjecture
- §3. Good Sections of Arithmetic Fundamental Groups
- §4. The Cuspidalisation of Sections of Arithmetic Fundamental groups
- §5. Applications to the Grothendieck Anabelian Section Conjecture
- §6. On a Weak Form of the p -adic Grothendieck Anabelian Section Conjecture

§0. Introduction. This, mostly expository, paper is around the topic of the Grothendieck anabelian section conjecture.

This conjecture predicts that splittings, or sections, of the exact sequence of the arithmetic fundamental group

$$1 \rightarrow \pi_1(\overline{X}) \rightarrow \pi_1(X) \rightarrow G_k \rightarrow 1$$

of a proper, smooth, and hyperbolic curve X , all arise from decomposition subgroups associated to rational points of X , over a base field k which is finitely generated over the prime field \mathbb{Q} (cf. §2).

A birational version of this conjecture predicts that splittings of the exact sequence of absolute Galois groups

$$1 \rightarrow \mathrm{Gal}(K_X^{\mathrm{sep}}/\overline{k}.K_X) \rightarrow \mathrm{Gal}(K_X^{\mathrm{sep}}/K_X) \rightarrow G_k \rightarrow 1,$$

where K_X is the function field of the hyperbolic curve X , all arise from rational points of X in some precise way (cf. §2), under the above assumption on the base field k (cf. §2).

This conjecture is one of the main topics in anabelian geometry. It establishes a dictionary between profinite group theory, arithmetic geometry, and Diophantine geometry.

This conjecture is still widely open. Only some examples are treated in the literature.

The most complete achievement around this conjecture, is the proof by Koenigsmann, and Pop, that the birational version of this conjecture holds true over p -adic local fields (cf. §5).

The main issue in investigating the section conjecture is the following. How can one produce a rational point $x \in X(k)$, from a splitting of the exact sequence of $\pi_1(X)$?

This issue is completely settled in the case of birational sections over p -adic local fields by Koenigsmann, and Pop. In this case, one produces a rational point by resorting to a local-global principle for Brauer groups of fields of transcendence degree 1 over p -adic local fields, that was proven by Lichtenbaum, in the finitely generated case, and by Pop in general.

It is not clear for the time being, at least to the author, how to settle the above issue in the case of sections of $\pi_1(X)$.

The main idea we would like to advocate in this paper is to reduce the solution of the section conjecture to the solution of its birational version.

We introduce, and investigate, the theory of cuspidalisation of sections of arithmetic fundamental groups for this purpose (cf. §4).

The main aim of this theory is, starting from a section

$$s : G_k \rightarrow \pi_1(X)$$

of the exact sequence of $\pi_1(X)$, to construct a section

$$\tilde{s} : G_k \rightarrow \text{Gal}(K_X^{\text{sep}}/K_X)$$

of the exact sequence of $\text{Gal}(K_X^{\text{sep}}/K_X)$, which lifts the section s , i.e. which inserts into a commutative diagram:

$$\begin{array}{ccc} G_k & \xrightarrow{\tilde{s}} & \text{Gal}(K_X^{\text{sep}}/K_X) \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s} & \pi_1(X) \end{array}$$

where the right vertical homomorphism is the natural one.

We have a natural exact sequence

$$1 \rightarrow I_X \rightarrow \text{Gal}(K_X^{\text{sep}}/K_X) \rightarrow \pi_1(X) \rightarrow 1,$$

where I_X is the normal subgroup of $\text{Gal}(K_X^{\text{sep}}/K_X)$ generated by the inertia subgroups at all geometric points of X .

We exhibit the notion of (uniformly) good sections of arithmetic fundamental groups (cf. §3). Sections which arise from rational points are good sections.

Our main result on the theory of cuspidalisation is that good sections $s : G_k \rightarrow \pi_1(X)$ of $\pi_1(X)$ can be lifted to sections

$$\tilde{s}^{c-ab} : G_k \rightarrow \text{Gal}(K_X^{\text{sep}}/K_X)^{c-ab}$$

of the natural projection $\mathrm{Gal}(K_X^{\mathrm{sep}}/K_X)^{c-\mathrm{ab}} \twoheadrightarrow G_k$, where $\mathrm{Gal}(K_X^{\mathrm{sep}}/K_X)^{c-\mathrm{ab}}$ is defined by the following push out diagram:

$$\begin{array}{ccccccc} 1 & \longrightarrow & I_X & \longrightarrow & \mathrm{Gal}(K_X^{\mathrm{sep}}/K_X) & \longrightarrow & \pi_1(X) \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \mathrm{id} \downarrow \\ 1 & \longrightarrow & I_X^{\mathrm{ab}} & \longrightarrow & \mathrm{Gal}(K_X^{\mathrm{sep}}/K_X)^{c-\mathrm{ab}} & \longrightarrow & \pi_1(X) \longrightarrow 1 \end{array}$$

and I_X^{ab} is the maximal abelian quotient of I_X , under quite general assumptions on the field k , which are satisfied by number fields, and p -adic local fields (cf. Corollary 4.7).

As an application, we prove a (pro- p) version of the section conjecture over p -adic local fields, for good sections of arithmetic fundamental groups, under some additional assumptions (cf. Theorem 5.4).

We state a (an unconditional) result on a semi-birational version of the p -adic section conjecture (cf. Theorem 5.6).

We also prove that the existence of good sections of $\pi_1(X)$, over number fields, implies the existence of degree 1 divisor on X , under a finiteness assumption of the Tate-Shafarevich group of the jacobian of X (cf. Theorem 5.8).

Finally, in §6, we discuss a weak version of the section conjecture over p -adic local fields, which is related to the absolute anabelian geometry of hyperbolic curves over p -adic local fields.

§1. Generalities on Arithmetic Fundamental Groups and Sections. In this section we introduce the set-up of the Grothendieck anabelian section conjecture.

1.1. Let k be a field of characteristic $p \geq 0$, and X a proper, smooth, geometrically connected, and hyperbolic algebraic curve over k . Let $K \stackrel{\mathrm{def}}{=} K_X$ be the function field of X , and η a geometric point of X above the generic point of X . Then η determines naturally an algebraic closure \bar{k} of k , a separable closure K_X^{sep} of K_X , and a geometric point $\bar{\eta}$ of $\bar{X} \stackrel{\mathrm{def}}{=} X \times_k \bar{k}$.

There exists a canonical exact sequence of profinite groups

$$(1.1) \quad 1 \rightarrow \pi_1(\bar{X}, \bar{\eta}) \rightarrow \pi_1(X, \eta) \xrightarrow{\mathrm{pr}_X} G_k \rightarrow 1.$$

Here, $\pi_1(X, \eta)$ denotes the arithmetic étale fundamental group of X with base point η , $\pi_1(\bar{X}, \bar{\eta})$ the étale fundamental group of \bar{X} with base point $\bar{\eta}$, and $G_k \stackrel{\mathrm{def}}{=} \mathrm{Gal}(\bar{k}/k)$ the absolute Galois group of k .

We will consider the following variant of the above exact sequence (1.1). Let

$$\Sigma \subseteq \mathfrak{Primes}$$

be a non-empty subset of the set \mathfrak{Primes} of all prime integers. In the case where $\mathrm{char}(k) = p > 0$, we will assume that $p \notin \Sigma$.

Write

$$\Delta_X \stackrel{\mathrm{def}}{=} \pi_1(\bar{X}, \bar{\eta})^\Sigma$$

for the maximal pro- Σ quotient of $\pi_1(\overline{X}, \overline{\eta})$, and

$$\Pi_X \stackrel{\text{def}}{=} \pi_1(X, \eta) / \text{Ker}(\pi_1(\overline{X}, \overline{\eta}) \twoheadrightarrow \pi_1(\overline{X}, \overline{\eta})^\Sigma)$$

for the quotient of $\pi_1(X, \eta)$ by the kernel of the natural surjective homomorphism $\pi_1(\overline{X}, \overline{\eta}) \twoheadrightarrow \pi_1(\overline{X}, \overline{\eta})^\Sigma$, which is a normal subgroup of $\pi_1(X, \eta)$.

Thus, we have an exact sequence of profinite groups

$$(1.2) \quad 1 \rightarrow \Delta_X \rightarrow \Pi_X \xrightarrow{\text{pr}_{X, \Sigma}} G_k \rightarrow 1.$$

We shall refer to $\pi_1(X, \eta)^{(\Sigma)} \stackrel{\text{def}}{=} \Pi_X$ as the geometrically pro- Σ quotient of $\pi_1(X, \eta)$, or the geometrically pro- Σ arithmetic fundamental group of X .

The exact sequence (1.2) induces a natural homomorphism

$$\rho_{X, \Sigma} : G_k \rightarrow \text{Out}(\Delta_X),$$

where $\text{Out}(\Delta_X) \stackrel{\text{def}}{=} \text{Aut}(\Delta_X) / \text{Inn}(\Delta_X)$ is the group of outer automorphisms of Δ_X . For $g \in G_k$, its image $\rho_{X, \Sigma}(g)$ in $\text{Out}(\Delta_X)$ is the class of the automorphism of Δ_X obtained by lifting g to an element $\tilde{g} \in \Pi_X$, and let \tilde{g} act on Δ_X by inner conjugation.

A profinite group G is slim if every open subgroup of G is centre-free (cf. [Mochizuki], §0). If G is a slim profinite group, we have a natural exact sequence

$$1 \rightarrow G \rightarrow \text{Aut } G \rightarrow \text{Out } G \rightarrow 1,$$

where the homomorphism $G \rightarrow \text{Aut } G$ sends an element $g \in G$ to the corresponding inner automorphism $h \mapsto ghg^{-1}$.

Moreover, if the profinite group G is finitely generated, then the groups $\text{Aut}(G)$, and $\text{Out}(G)$, are naturally endowed with a profinite topology, and the above sequence is an exact sequence of profinite groups.

The following is an important property of the profinite groups Δ_X , and Π_X .

Lemma 1.2. *The profinite group Δ_X is slim. In particular, the exact sequence (1.2) is obtained from the following exact sequence*

$$(1.3) \quad 1 \rightarrow \Delta_X \rightarrow \text{Aut}(\Delta_X) \rightarrow \text{Out}(\Delta_X) \rightarrow 1,$$

by pull back via the natural continuous homomorphism

$$\rho_{X, \Sigma} : G_k \rightarrow \text{Out}(\Delta_X).$$

More precisely, we have a commutative diagram:

$$(1.4) \quad \begin{array}{ccccccc} 1 & \longrightarrow & \Delta_X & \longrightarrow & \text{Aut}(\Delta_X) & \longrightarrow & \text{Out}(\Delta_X) \longrightarrow 1 \\ & & \text{id} \uparrow & & \uparrow & & \rho_{X, \Sigma} \uparrow \\ 1 & \longrightarrow & \Delta_X & \longrightarrow & \Pi_X & \longrightarrow & G_k \longrightarrow 1 \end{array}$$

where the horizontal arrows are exact, and the right square is cartesian.

Proof. Well known, see for example [Tamagawa], Proposition 1.11. \square

1.3. Similarly, if we write $X \times X \stackrel{\text{def}}{=} X \times_k X$, and $\iota : X \rightarrow X \times X$ for the natural diagonal embedding, then the geometric point η determines naturally (via ι) a geometric point, which we will also denote η , of $X \times X$.

There exists a natural exact sequence of profinite groups

$$(1.5) \quad 1 \rightarrow \pi_1(\overline{X \times X}, \bar{\eta}) \rightarrow \pi_1(X \times X, \eta) \xrightarrow{\text{pr}} G_k \rightarrow 1.$$

Here, $\pi_1(X \times X, \eta)$ denotes the arithmetic étale fundamental group of $X \times X$ with base point η , which is naturally identified with the fibre product $\pi_1(X, \eta) \times_{G_k} \pi_1(X, \eta)$, and $\pi_1(\overline{X \times X}, \bar{\eta})$ is the étale fundamental group of $\overline{X \times X}$ with base point $\bar{\eta}$ ($\bar{\eta}$ is naturally induced by η), which is naturally identified with the product $\pi_1(\overline{X}, \bar{\eta}) \times \pi_1(\overline{X}, \bar{\eta})$.

Similarly, as in 1.1, we consider the maximal pro- Σ quotient

$$\Delta_{X \times X} \stackrel{\text{def}}{=} \pi_1(\overline{X \times X}, \bar{\eta})^\Sigma$$

of $\pi_1(\overline{X \times X}, \bar{\eta})$, which is naturally identified with $\Delta_X \times \Delta_X$, and the geometrically pro- Σ quotient

$$\Pi_{X \times X} \stackrel{\text{def}}{=} \pi_1(X \times X, \eta)^{(\Sigma)} \stackrel{\text{def}}{=} \pi_1(X \times X, \eta) / \text{Ker}(\pi_1(\overline{X \times X}, \bar{\eta}) \twoheadrightarrow \pi_1(\overline{X \times X}, \bar{\eta})^\Sigma)$$

of $\pi_1(X \times X, \eta)$, which is naturally identified with $\Pi_X \times_{G_k} \Pi_X$.

Thus, we have a natural exact sequence

$$1 \rightarrow \Delta_{X \times X} \rightarrow \Pi_{X \times X} \rightarrow G_k \rightarrow 1.$$

1.4. Our main objects of interest are group-theoretic splittings, or sections, of the exact sequence (1.2).

Let

$$s : G_k \rightarrow \Pi_X$$

be a continuous group-theoretic section of the natural projection

$$\text{pr} \stackrel{\text{def}}{=} \text{pr}_{X, \Sigma} : \Pi_X \twoheadrightarrow G_k,$$

meaning that $\text{pr} \circ s : G_k \rightarrow G_k$ is the identity homomorphism.

We will refer to $s : G_k \rightarrow \Pi_X$, as above, as a section of the arithmetic fundamental group Π_X .

Every inner automorphism $\text{inn}^g : \Pi_X \rightarrow \Pi_X$ of Π_X by an element $g \in \Delta_X$, gives rise to a conjugate section $\text{inn}^g \circ s : G_k \rightarrow \Pi_X$. We will refer to the set

$$\mathcal{C}[s] \stackrel{\text{def}}{=} \{\text{inn}^g \circ s : G_k \rightarrow \Pi_X\}_{g \in \Delta_X}$$

as the set of conjugacy classes of the section s .

§2. Grothendieck Anabelian Section Conjecture. We follow the same notations as in §1.

2.1. Sections of arithmetic fundamental groups arise naturally from rational points.

More precisely, Let $x \in X(k)$ be a rational point. Then x determines a decomposition subgroup

$$D_x \subset \Pi_X,$$

which is defined only up to conjugation by the elements of Δ_X , and which maps isomorphically to G_k via the projection $\text{pr} : \Pi_X \twoheadrightarrow G_k$.

Hence, the subgroup $D_x \subset \Pi_X$, determines a group-theoretic section

$$s_x : G_k \rightarrow \Pi_X$$

of the natural projection $\text{pr} : \Pi_X \twoheadrightarrow G_k$, which is defined only up to conjugation by the elements of Δ_X .

Let $\overline{\text{Sec}}_{\Pi_X}$ be the set of conjugacy classes of all continuous group-theoretic sections $G_k \rightarrow \Pi_X$, of the natural projection $\text{pr} : \Pi_X \twoheadrightarrow G_k$, modulo inner conjugation by the elements of Δ_X (cf. 1.4).

We have a natural set-theoretic map

$$\varphi_{X,\Sigma} : X(k) \rightarrow \overline{\text{Sec}}_{\Pi_X},$$

$$x \mapsto \varphi_{X,\Sigma}(x) \stackrel{\text{def}}{=} [s_x],$$

where $[s_x]$ denotes the class of a section $s_x : G_k \rightarrow \Pi_X$, associated to the rational point x , in $\overline{\text{Sec}}_{\Pi_X}$.

Definition 2.2. Let $s : G_k \rightarrow \Pi_X$ be a continuous group-theoretic section of the natural projection $\text{pr} : \Pi_X \twoheadrightarrow G_k$. We say that the section s is point-theoretic, or geometric, if the class $[s]$ of s in $\overline{\text{Sec}}_{\Pi_X}$ belongs to the image of the map $\varphi_{X,\Sigma}$.

The following is the main Grothendieck anabelian conjecture regarding sections of arithmetic fundamental groups. This conjecture establishes a dictionary between purely group-theoretic sections of arithmetic fundamental groups, and rational points of hyperbolic curves over finitely generated fields, in characteristic 0.

Grothendieck Anabelian Section Conjecture (GASC). (cf. [Grothendieck]) Assume that k is finitely generated over the prime field \mathbb{Q} , and $\Sigma = \mathfrak{Primes}$. Then the map $\varphi_X \stackrel{\text{def}}{=} \varphi_{X,\Sigma} : X(k) \rightarrow \overline{\text{Sec}}_{\Pi_X}$ is bijective. In particular, every group-theoretic section of Π_X is point-theoretic in this case.

The injectivity of the map φ_X is well-known (cf. for example [Mochizuki1], Theorem 19.1).

So the statement of the GASC is equivalent to the surjectivity of the set-theoretic map φ_X , i.e. that every group-theoretic section of Π_X is point-theoretic, under the above assumptions.

Note that a similar conjecture can be formulated over any field, and for any non-empty set Σ of prime integers, but one can not expect its validity in general.

For example, the analog of this conjecture doesn't hold over finite fields (even if $\Sigma = \mathfrak{Primes}$).

Indeed, over a finite field the natural projection $\Pi_X \twoheadrightarrow G_k$ admits group-theoretic sections, or splittings, since the profinite group G_k is free in this case. On the other hand, there are proper, smooth, geometrically connected, and hyperbolic curves over finite fields with no rational points.

2.3. Assume that k is a number field, i.e. k is a finite extension of the field of rational numbers \mathbb{Q} . Let v be a place of k , and denote by k_v the completion of k at v . Write

$$X_v \stackrel{\text{def}}{=} X \times_k k_v.$$

Let $D_v \subset G_k$ be a decomposition group at v (D_v is only defined up to conjugation), which is naturally isomorphic to the absolute Galois group G_{k_v} of k_v .

By pulling back the exact sequence

$$1 \rightarrow \Delta_X \rightarrow \Pi_X \rightarrow G_k \rightarrow 1,$$

by the natural injective homomorphism $D_v \hookrightarrow G_k$, we obtain the exact sequence

$$1 \rightarrow \Delta_{X_v} \rightarrow \Pi_{X_v} \rightarrow G_{k_v} \rightarrow 1.$$

Note that there exists an isomorphism $\Delta_X \xrightarrow{\sim} \Delta_{X_v}$.

In particular, a group-theoretic section $s : G_k \rightarrow \Pi_X$ of the arithmetic fundamental group Π_X , induces naturally a group-theoretic section

$$s_v : G_{k_v} \rightarrow \Pi_{X_v}$$

of Π_{X_v} , for each place v of k .

Moreover, if the section s is point-theoretic, then the section s_v is point-theoretic, for each place v of k , as is easily seen.

It seems quite natural to formulate an analog of the GASC over p -adic local fields.

A p -adic Version of Grothendieck Anabelian Section Conjecture (p-adic GASC). Assume that k is a p -adic local field, i.e. k is a finite extension of the field \mathbb{Q}_p , for some prime integer p , and $\Sigma = \mathfrak{P}\text{rim}\mathfrak{es}$. Then the map $\varphi_X \stackrel{\text{def}}{=} \varphi_X \stackrel{\text{def}}{=} \varphi_{X,\Sigma} : X(k) \rightarrow \overline{\text{Sec}}_{\Pi_X}$ is bijective.

The map φ_X is known to be injective in the case where k is a p -adic local field, and $p \in \Sigma$ (cf. [Mochizuki1], Theorem 19.1). Thus, the statement of the p -adic GASC is equivalent to the surjectivity of the map φ_X in this case.

2.4. Next, we recall the definition of a system of neighbourhoods of a group-theoretic section of the arithmetic fundamental group Π_X .

The profinite group Δ_X being topologically finitely generated, there exists a sequence of characteristic open subgroups

$$\dots \subseteq \Delta_X[i+1] \subseteq \Delta_X[i] \subseteq \dots \subseteq \Delta_X[1] \stackrel{\text{def}}{=} \Delta_X$$

(where i ranges over all positive integers) of Δ_X , such that

$$\bigcap_{i \geq 1} \Delta_X[i] = \{1\}.$$

In particular, given a group-theoretic section $s : G_k \rightarrow \Pi_X$ of Π_X , we obtain open subgroups

$$\Pi_X[i, s] \stackrel{\text{def}}{=} s(G_k) \cdot \Delta_X[i] \subseteq \Pi_X$$

(where $s(G_k)$ denotes the image of G_k in Π_X via the section s) of Π_X , whose intersection coincide with $s(G_k)$, and which correspond to a tower of finite étale (not necessarily Galois) covers

$$\dots \rightarrow X_{i+1}[s] \rightarrow X_i[s] \rightarrow \dots \rightarrow X_1[s] \stackrel{\text{def}}{=} X$$

defined over k .

We will refer to the set $\{X_i[s]\}_{i \geq 1}$ as a system of neighbourhoods of the section s .

Note that for each positive integer i , the open subgroup $\Pi_X[i, s]$ of Π_X is naturally identified with the geometrically pro- Σ arithmetic étale fundamental group $\pi_1(X_i[s], \eta_i)^{(\Sigma)}$ (the geometric point η_i of $X_i[s]$ is naturally induced by the geometric point η of X), and sits naturally in the following exact sequence

$$1 \rightarrow \Delta_X[i] \rightarrow \Pi_X[i, s] \rightarrow G_k \rightarrow 1,$$

which inserts in the following commutative diagram:

$$\begin{array}{ccccccc} 1 & \longrightarrow & \Delta_X[i] & \longrightarrow & \Pi_X[i, s] & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \text{id} \downarrow \\ 1 & \longrightarrow & \Delta_X & \longrightarrow & \Pi_X & \longrightarrow & G_k \longrightarrow 1 \end{array}$$

where the two left vertical homomorphisms are the natural inclusions.

In particular, by the very definition of $\Pi_X[i, s]$, the section s restricts naturally to a group-theoretic section

$$s_i : G_k \rightarrow \Pi_X[i, s]$$

of the natural projection $\Pi_X[i, s] \twoheadrightarrow G_k$, which fits into the following commutative diagram:

$$\begin{array}{ccc} G_k & \xrightarrow{s_i} & \Pi_X[i, s] \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s} & \Pi_X \end{array}$$

where the right vertical homomorphism is the natural inclusion.

Thus, a section $s : G_k \rightarrow \Pi_X$ gives rise to a system of neighbourhoods $\{X_i[s]\}_{i \geq 1}$, and the corresponding open subgroups $\{\Pi_X[i, s]\}_{i \geq 1}$ inherit naturally, from the section s , sections $s_i : G_k \rightarrow \Pi_X[i, s]$.

In investigating the point-theorecity of the section s (cf. Definition 2.2), it is important to observe not only the section s , but rather the family of sections $\{s_i\}_{i \geq 1}$.

In fact, a number of important properties of the section s can be proven, using a limit argument, by observing the family of sections $\{s_i\}_{i \geq 1}$, rather than only the section s .

The best illustration of this phenomenon is the following crucial observation, which is extremely important in investigating the Grothendieck anabelian section conjecture, and which is due to Tamagawa.

Lemma 2.5. *Assume that k is finitely generated over the prime field \mathbb{Q} , or that k is a p -adic local field. Let $s : G_k \rightarrow \Pi_X$ be a group-theoretic section of the natural projection $\text{pr} : \Pi_X \twoheadrightarrow G_k$, and $\{X_i[s]\}_{i \geq 1}$ a system of neighbourhoods of the section s (cf. 2.4). Then the section s is point-theoretic if and only if $X_i[s](k) \neq \emptyset$, for each $i \geq 1$.*

Proof. See [Tamagawa], Proposition 2.8, (iv). \square

The above Lemma 2.5 reduces the proof of the Grothendieck anabelian section conjecture, in the case where k is finitely generated over the prime field \mathbb{Q} , or that k is a p -adic local field, and $\Sigma = \mathfrak{Primes}$, to proving the following implication

$$\{\overline{\text{Sec}}_{\Pi_X} \neq \emptyset\} \implies \{X(k) \neq \emptyset\}.$$

Thus, at the heart of the Grothendieck anabelian section conjecture, is the following fundamental problem.

Problem. How can one produce, under the assumptions of GASC, or the p -adic GASC, a rational point $x \in X(k)$ starting from a group-theoretic section $s : G_k \rightarrow \Pi_X$ of Π_X ?

No systematic approach has yet been developed so far to attack this problem.

The only situation where a method, or a technique, is available to solve this problem positively is the method, developed by Koenigsmann, and Pop, in the framework of the birational version of the p -adic GASC (cf. [Koenigsmann], and [Pop]), and which resorts to a local-global principle for Brauer groups of fields of transcendence degree 1 over p -adic local fields (see the discussion after Theorem 5.2).

Remarks 2.6.

(i) One of the difficulties in investigating the GASC is that, for the time being, one doesn't know how to construct sections of arithmetic fundamental groups, and hence test the validity of the conjecture on concrete examples.

In fact, the GASC itself can be viewed as a "rigidity" statement.

Namely, the only way one knows (so far) to construct sections of arithmetic fundamental groups of hyperbolic curves, over finitely generated fields of characteristic 0, is via decomposition groups associated to rational points, and these should be the only sections that exist !

One way, however, to construct such sections is as follows.

Let X be a proper, smooth, geometrically connected, hyperbolic algebraic curve over the field k , and $\Sigma \subseteq \mathfrak{Primes}$ a non-empty set of prime integers.

Consider the exact sequence

$$(1.2) \quad 1 \rightarrow \Delta_X \rightarrow \Pi_X \xrightarrow{\text{pr}} G_k \rightarrow 1,$$

where Π_X is the geometrically pro- Σ arithmetic fundamental group of X .

Recall the commutative cartesian diagram:

$$(1.4) \quad \begin{array}{ccccccc} 1 & \longrightarrow & \Delta_X & \longrightarrow & \text{Aut}(\Delta_X) & \longrightarrow & \text{Out}(\Delta_X) \longrightarrow 1 \\ & & \uparrow \text{id} & & \uparrow & & \uparrow \rho_{X, \Sigma} \\ 1 & \longrightarrow & \Delta_X & \longrightarrow & \Pi_X & \xrightarrow{\text{pr}} & G_k \longrightarrow 1 \end{array}$$

In order to construct a continuous group-theoretic section $s : G_k \rightarrow \Pi_X$ of the natural projection $\text{pr} : \Pi_X \twoheadrightarrow G_k$, it is equivalent to construct a continuous homomorphism $\tilde{\rho}_{X,\Sigma} : G_k \rightarrow \text{Aut}(\Delta_X)$, which lifts the homomorphism $\rho_{X,\Sigma} : G_k \rightarrow \text{Out}(\Delta_X)$ above, i.e. such that the following diagram commutes

$$\begin{array}{ccc} G_k & \xrightarrow{\tilde{\rho}_{X,\Sigma}} & \text{Aut}(\Delta_X) \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{\rho_{X,\Sigma}} & \text{Out}(\Delta_X) \end{array}$$

as follows directly from the fact that the right square in the diagram (1.4) is cartesian.

(ii) In light of the Remark (i), it is possible to construct sections $s : G_k \rightarrow \Pi_X$ of Π_X , if the image of G_k in $\text{Out}(\Delta_X)$ via the natural homomorphism $\rho_{X,\Sigma} : G_k \rightarrow \text{Out}(\Delta_X)$ is a free pro- Σ group.

Under this condition one may hope to construct non geometric sections s , i.e. sections which do not arise from rational points. This is the method used in [Hoshi] to construct non geometric sections in the case where k is a number field, or a p -adic local field, and $\Sigma = \{p\}$.

However, this method is unlikely to produce examples of non-geometric sections in the case where $\Sigma = \mathfrak{Primes}$. Indeed, in this case the homomorphism $\rho_{X,\Sigma} : G_k \rightarrow \text{Out}(\Delta_X)$ is injective if X is hyperbolic, and k is a number field, or a p -adic local field, as is well-known.

2.7. One can formulate a birational version of the Grothendieck anabelian section conjecture as follows (see also [Pop]).

There exists a natural exact sequence of absolute Galois groups

$$1 \rightarrow \text{Gal}(K_X^{\text{sep}}/K_X.\bar{k}) \rightarrow \text{Gal}(K_X^{\text{sep}}/K_X) \rightarrow G_k \rightarrow 1.$$

Let $\Sigma \subseteq \mathfrak{Primes}$ be a non-empty set of prime integers, and

$$\overline{G}_X \stackrel{\text{def}}{=} \text{Gal}(K_X^{\text{sep}}/K_X.\bar{k})^\Sigma$$

the maximal pro- Σ quotient of the absolute Galois group $\text{Gal}(K_X^{\text{sep}}/K_X.\bar{k})$.

Let

$$G_X \stackrel{\text{def}}{=} \text{Gal}(K_X^{\text{sep}}/K_X) / \text{Ker}(\text{Gal}(K_X^{\text{sep}}/K_X.\bar{k}) \twoheadrightarrow \text{Gal}(K_X^{\text{sep}}/K_X.\bar{k})^\Sigma)$$

be the maximal geometrically pro- Σ Galois group of the function field K_X .

Thus, G_X sits naturally in the following exact sequence

$$1 \rightarrow \overline{G}_X \rightarrow G_X \rightarrow G_k \rightarrow 1.$$

Let $x \in X(k)$ be a rational point. Then x determines a decomposition subgroup $D_x \subset G_X$, which is only defined up to conjugation by the elements of \overline{G}_X , and which maps onto G_k via the natural projection $G_X \twoheadrightarrow G_k$.

More precisely, D_x sits naturally in the following exact sequence

$$1 \rightarrow M_X \rightarrow D_x \rightarrow G_k \rightarrow 1,$$

where $M_X \xrightarrow{\sim} \hat{\mathbb{Z}}(1)^\Sigma$ (cf. 3.1, for the definition of the module of roots of unity M_X).

The above sequence is known to be split. The set of all possible splittings, i.e. sections $G_k \rightarrow D_x$ of the above exact sequence, is a torsor under $H^1(G_k, M_X)$. The latter can be naturally identified, via Kummer theory, with the Σ -adic completion $\hat{k}^{\times, \Sigma}$ of the multiplicative group k^\times .

Each section $G_k \rightarrow D_x$ of the natural projection $D_x \twoheadrightarrow G_k$ determines naturally a section $G_k \rightarrow G_X$ of the natural projection $G_X \twoheadrightarrow G_k$, whose image is contained in D_x .

The Birational Grothendieck Anabelian Section Conjecture (BGASC).

Assume that k is finitely generated over the prime field \mathbb{Q} , and $\Sigma = \mathfrak{Primes}$. Let $s : G_k \rightarrow G_X$ be a group-theoretic section of the natural projection $G_X \twoheadrightarrow G_k$. Then the image $s(G_k)$ is contained in a decomposition subgroup $D_x \subset G_X$ associated to a unique rational point $x \in X(k)$. In particular, the existence of the section s implies that $X(k) \neq \emptyset$.

One can, in a similar way, formulate a p -adic version of this conjecture.

A p -adic Version of The Birational Grothendieck Anabelian Section Conjecture (p -adic BGASC).

Assume that k is a p -adic local field, i.e. k is a finite extension of \mathbb{Q}_p , and $\Sigma = \mathfrak{Primes}$. Let $s : G_k \rightarrow G_X$ be a group-theoretic section of the natural projection $G_X \twoheadrightarrow G_k$. Then the image $s(G_k)$ is contained in a decomposition subgroup $D_x \subset G_X$ associated to a unique rational point $x \in X(k)$. In particular, the existence of the section s implies that $X(k) \neq \emptyset$.

§3. Good Sections of Arithmetic Fundamental Groups. We use the same notations as in §1, and §2.

We will introduce the notion of (uniformly) good sections of arithmetic fundamental groups. Point-theoretic sections of arithmetic fundamental groups (cf. Definition 2.2) are (uniformly) good sections.

3.1. Next, we recall the definition of the arithmetic Chern class associated to a group-theoretic section s of the arithmetic fundamental group Π_X (cf. [Saïdi], 1.2, and [Esnault-Wittenberg], for more details).

In what follows all scheme cohomology groups are étale cohomology groups.

First, let $\hat{\mathbb{Z}}^\Sigma$ be the maximal pro- Σ quotient of $\hat{\mathbb{Z}}$, and

$$M_X \stackrel{\text{def}}{=} \text{Hom}(\mathbb{Q}/\mathbb{Z}, (K_X^{\text{sep}})^\times) \otimes_{\hat{\mathbb{Z}}} \hat{\mathbb{Z}}^\Sigma.$$

Note that M_X is a free $\hat{\mathbb{Z}}^\Sigma$ -module of rank one, and has a natural structure of G_k -module, which is isomorphic to the G_k -module $\hat{\mathbb{Z}}(1)^\Sigma$, where the “(1)” denotes a Tate twist, i.e. G_k acts on $\hat{\mathbb{Z}}(1)^\Sigma$ via the Σ -part of the cyclotomic character. We will refer to M_X as the module of roots of unity attached to X , relative to the set of primes Σ .

Let

$$\eta_X^{\text{diag}} \in H^2(X \times X, M_X)$$

be the étale Chern class, which is associated to the diagonal embedding $\iota : X \rightarrow X \times_k X$, or alternatively the first Chern class of the line bundle $\mathcal{O}_{X \times X}(\iota(X))$.

There exists a natural identification (cf. [Mochizuki], Proposition 1.1)

$$H^2(X \times X, M_X) \xrightarrow{\sim} H^2(\Pi_{X \times X}, M_X).$$

The Chern class η_X^{diag} corresponds via the above identification to an extension class

$$\eta_X^{\text{diag}} \in H^2(\Pi_{X \times X}, M_X).$$

We shall refer to the extension class η_X^{diag} as the extension class of the diagonal.

Let $s : G_k \rightarrow \Pi_X$ be a group-theoretic section of the natural projection $\Pi_X \rightarrow G_k$. Let

$$(3.1) \quad 1 \rightarrow M_X \rightarrow \mathcal{D} \rightarrow \Pi_{X \times X} \rightarrow 1$$

be a group extension, whose class in $H^2(\Pi_{X \times X}, M_X)$ coincides with the extension class η_X^{diag} of the diagonal.

By pulling back the group extension (3.1) by the continuous injective homomorphism

$$(s, \text{id}) : G_k \times_{G_k} \Pi_X \rightarrow \Pi_{X \times X},$$

we obtain a natural commutative diagram:

$$\begin{array}{ccccccc} 1 & \longrightarrow & M_X & \longrightarrow & \mathcal{D}_s & \longrightarrow & G_k \times_{G_k} \Pi_X \longrightarrow 1 \\ & & \text{id} \downarrow & & \downarrow & & (s, \text{id}) \downarrow \\ 1 & \longrightarrow & M_X & \longrightarrow & \mathcal{D} & \longrightarrow & \Pi_{X \times X} \longrightarrow 1 \end{array}$$

where the right square is cartesian.

Further, via the natural identification $G_k \times_{G_k} \Pi_X \xrightarrow{\sim} \Pi_X$, the upper group extension \mathcal{D}_s in the above diagram corresponds to a group extension (which we denote also \mathcal{D}_s)

$$1 \rightarrow M_X \rightarrow \mathcal{D}_s \rightarrow \Pi_X \rightarrow 1.$$

We will refer to the class $[\mathcal{D}_s]$ of the extension \mathcal{D}_s in $H^2(\Pi_X, M_X)$ as the extension class associated to the section s .

Definition 3.2 (The (Σ) -Étale Chern Class associated to a Section). We define the (Σ) -étale Chern class $c(s) \in H^2(X, M_X)$ associated to the section s as the element of $H^2(X, M_X)$ corresponding to the above extension class $[\mathcal{D}_s]$, which is associated to the section s , via the natural identification $H^2(\Pi_X, M_X) \xrightarrow{\sim} H^2(X, M_X)$.

Let $\{X_i[s]\}_{i \geq 1}$ be a system of neighbourhoods of the section s , and $\{\Pi_X[i, s]\}_{i \geq 1}$ the corresponding open subgroups of Π_X (cf. 2.4). Recall that the section s restricts naturally to a group-theoretic section

$$s_i : G_k \rightarrow \Pi_X[i, s]$$

of the natural projection $\Pi_X[i, s] \twoheadrightarrow G_k$, which fits into the following commutative diagram:

$$\begin{array}{ccc} G_k & \xrightarrow{s_i} & \Pi_X[i, s] \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s} & \Pi_X \end{array}$$

where the right vertical homomorphism is the natural inclusion, for each positive integer i (cf. loc. cit).

One can easily observe the following Lemma (See [Saïdi], Lemma 1.3.1, for more details).

Lemma 3.3. *For each positive integer i , the image of the Chern class $c(s_{i+1}) \in H^2(X_{i+1}[s], M_X)$ associated to the section $s_{i+1} : G_k \rightarrow \Pi_X[i+1, s]$ in $H^2(X[i, s], M_X)$, via the corestriction homomorphism $\text{cor} : H^2(X_{i+1}[s], M_X) \rightarrow H^2(X_i[s], M_X)$, coincides with the Chern class $c(s_i)$ associated to the section $s_i : G_k \rightarrow \Pi_X[i, s]$.*

Definition 3.4 (The pro-(Σ)-Étale Chern Class associated to a Section). Let $\varprojlim_{i \geq 1} H^2(X_i[s], M_X)$ be the projective limit of the $H^2(X_i[s], M_X)$'s, where the transition homomorphisms are the corestriction homomorphisms. We define the pro-(Σ)-étale Chern class associated to the section s , relative to the system of neighbourhoods $\{X_i[s]\}_{i \geq 1}$, as the element $\hat{c}(s) \stackrel{\text{def}}{=} (c(s_i))_{i \geq 1} \in \varprojlim_{i \geq 1} H^2(X_i[s], M_X)$ (cf. Lemma 3.3).

3.5. Next, we will introduce the notion of (uniformly) good sections of arithmetic fundamental groups.

For each positive Σ -integer n , meaning that n is an integer which is divisible only by primes in Σ , the Kummer exact sequence in étale topology

$$1 \rightarrow \mu_n \rightarrow \mathbb{G}_m \xrightarrow{n} \mathbb{G}_m \rightarrow 1$$

induces naturally, for each positive integer i , an exact sequence of abelian groups

$$(3.2) \quad 0 \rightarrow \text{Pic}(X_i[s])/n \text{Pic}(X_i[s]) \rightarrow H^2(X_i[s], \mu_n) \rightarrow_n \text{Br}(X_i[s]) \rightarrow 0,$$

which for positive integers m and n , with n divides m , fits naturally into a commutative diagram:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \text{Pic}(X_i[s])/m \text{Pic}(X_i[s]) & \longrightarrow & H^2(X_i[s], \mu_m) & \longrightarrow & {}_m \text{Br}(X_i[s]) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \text{Pic}(X_i[s])/n \text{Pic}(X_i[s]) & \longrightarrow & H^2(X_i[s], \mu_n) & \longrightarrow & {}_n \text{Br}(X_i[s]) & \longrightarrow & 0 \end{array}$$

where the lower and upper horizontal sequences are the above exact sequence (3.2), and the vertical homomorphisms are the natural homomorphisms.

Here, $\text{Pic} \stackrel{\text{def}}{=} H^1(\cdot, \mathbb{G}_m)$ denotes the Picard group, $\text{Br} \stackrel{\text{def}}{=} H^2(\cdot, \mathbb{G}_m)$ is the Brauer-Grothendieck cohomological group, and for a positive integer n : ${}_n \text{Br} \subseteq \text{Br}$ is the subgroup of Br which is annihilated by n .

By taking projective limits, the above diagram induces naturally, for each positive integer i , the following exact sequence

$$0 \rightarrow \varprojlim_{n \text{ } \Sigma\text{-integer}} \text{Pic}(X_i[s])/n \text{Pic}(X_i[s]) \rightarrow H^2(X_i[s], M_X) \rightarrow \varprojlim_{n \text{ } \Sigma\text{-integer}} {}_n \text{Br}(X_i[s]) \rightarrow 0.$$

We will denote by

$$\text{Pic}(X_i[s])^{\wedge, \Sigma} \stackrel{\text{def}}{=} \varprojlim_{n \text{ } \Sigma\text{-integer}} \text{Pic}(X_i[s])/n \text{Pic}(X_i[s])$$

the Σ -adic completion of the Picard group $\text{Pic}(X_i[s])$, and

$$T \text{Br}(X_i[s])^{\Sigma} \stackrel{\text{def}}{=} \varprojlim_{n \text{ } \Sigma\text{-integer}} {}_n \text{Br}(X_i[s])$$

the Σ -Tate module of the Brauer group $\text{Br}(X_i[s])$. Thus, we have a natural exact sequence

$$(3.3) \quad 0 \rightarrow \text{Pic}(X_i[s])^{\wedge, \Sigma} \rightarrow H^2(X_i[s], M_X) \rightarrow T\text{Br}(X_i[s])^{\Sigma} \rightarrow 0.$$

In what follows we will identify $\text{Pic}(X_i[s])^{\wedge, \Sigma}$ with its image in $H^2(X_i[s], M_X)$ (cf. the above exact sequence (3.3)), and refer to it as the Picard part of $H^2(X_i[s], M_X)$.

Let

$$s : G_k \rightarrow \Pi_X[1, s] \stackrel{\text{def}}{=} \Pi_X$$

be a group-theoretic section of Π_X as above. For each positive integer i , let

$$s_i : G_k \rightarrow \Pi_X[i, s]$$

be the induced group-theoretic section of $\Pi_X[i, s]$.

By pulling back cohomology classes via the section s_i , and bearing in mind the natural identifications $H^2(\Pi_X[i, s], M_X) \xrightarrow{\sim} H^2(X_i[s], M_X)$ (cf. [Mochizuki, Proposition 1.1]), we obtain a natural (restriction) homomorphism

$$s_i^* : H^2(X_i[s], M_X) \rightarrow H^2(G_k, M_X).$$

Finally, observe that if k' is a finite extension of k , and $X_{k'} \stackrel{\text{def}}{=} X \times_k k'$, then we have a natural commutative diagram:

$$\begin{array}{ccccccc} 1 & \longrightarrow & \Delta_X & \longrightarrow & \Pi_{X_{k'}} \stackrel{\text{def}}{=} \pi_1(X_{k'}, \eta)^{(\Sigma)} & \longrightarrow & G_{k'} \longrightarrow 1 \\ & & \text{id} \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \Delta_X & \longrightarrow & \Pi_X & \longrightarrow & G_k \longrightarrow 1 \end{array}$$

where the right, and middle, vertical arrows are the natural inclusions, and the far right square is cartesian.

In particular, the section $s_k \stackrel{\text{def}}{=} s : G_k \rightarrow \Pi_X$ induces naturally a group-theoretic section $s_{k'} : G_{k'} \rightarrow \Pi_{X_{k'}}$ of $\Pi_{X_{k'}}$.

Definition 3.6 (Good and Uniformly Good Sections of Arithmetic Fundamental Groups). We say that the section s is a good group-theoretic section, relative to the system of neighbourhoods $\{X_i[s]\}_{i \geq 1}$, if the above homomorphisms $s_i^* : H^2(X_i[s], M_X) \rightarrow H^2(G_k, M_X)$, for each positive integer $i \geq 1$, annihilate the Picard part $\text{Pic}(X_i[s])^{\wedge, \Sigma}$ of $H^2(X_i[s], M_X)$. In other words, the section s is good if $\text{Pic}(X_i[s])^{\wedge, \Sigma} \subseteq \text{Ker } s_i^*$, for each positive integer i .

We say that the section s is uniformly good, relative to the system of neighbourhoods $\{X_i[s]\}_{i \geq 1}$, if the induced section $s_{k'} : G_{k'} \rightarrow \Pi_{X_{k'}}$ is good, relative to the system of neighbourhoods of $s_{k'}$ which is naturally induced by the $\{X_i[s]\}_{i \geq 1}$, for every finite extension k'/k .

It is easy to see that the above definition is independent of the given system of neighbourhoods $\{X_i[s]\}_{i \geq 1}$ of the section s . We will refer to a section satisfying the conditions in Definition 3.6 as good, or uniformly good, without necessarily specifying a system of neighbourhoods of the section.

The notion of uniformly good sections is motivated by the fact that a necessary condition for a group-theoretic section $s : G_k \rightarrow \Pi_X$ to be point-theoretic, i.e. arises from a k -rational point $x \in X(k)$ (cf. Definition 2.2), is that the section s is uniformly good in the sense of Definition 3.6 (cf. [Saïdi, Proposition 1.5.2]).

Remarks 3.7.

(i) If k is a p -adic local field, the conditions of goodness and uniform goodness for the section s are equivalent.

Moreover, if $p \in \Sigma$, the section s is good in this case if and only if $X(k^{\text{tame}}) \neq \emptyset$, where k^{tame} is the maximal tame extension of k (cf. [Saïdi], Proposition 1.6.6, and Proposition 1.6.8).

(ii) If k is a number field, one has a local-global principle for (uniform) goodness. Namely, the section s is good in this case if and only if the section s_v is good, for each place v of k (cf. loc. cit., Proposition 1.8.1).

Remark 3.8. Another necessary condition for the section s to be point-theoretic, is that the image of the pro-Chern class $\hat{c}(s) \in \varprojlim_{i \geq 1} H^2(X_i[s], M_X)$ in $\varprojlim_{i \geq 1} T \text{Br}(X_i[s])^\Sigma$, via the natural homomorphism

$$\varprojlim_{i \geq 1} H^2(X_i[s], M_X) \rightarrow \varprojlim_{i \geq 1} T \text{Br}(X_i[s])^\Sigma,$$

equals 0.

In other words, if the section s is point-theoretic, then the associated pro-Chern class $\hat{c}(s)$ (cf. Definition 3.4) lies in the Picard part $\varprojlim_{i \geq 1} \text{Pic}(X_i[s])^{\wedge, \Sigma}$ of

$$\varprojlim_{i \geq 1} H^2(X_i[s], M_X).$$

Indeed, Assume that the section s is point-theoretic, meaning that $s \stackrel{\text{def}}{=} s_x : G_k \rightarrow \Pi_X$ arises from a k -rational point $x \in X(k)$ (cf. Definition 2.2). Then there exists a compatible system of rational points $\{x_i \in X_i[s](k)\}_{i \geq 1}$, i.e. x_{i+1} maps to x_i via the natural morphism $X_{i+1}[s] \rightarrow X_i[s]$.

For every positive integer i , let $\mathcal{O}(x_i) \in \text{Pic}(X_i[s])$ be the degree 1 line bundle associated to x_i .

The Chern class $c(s_i) \in H^2(X_i[s], M_X)$, which is associated to the section s_i (cf. Definition 3.2), coincides with the étale Chern class $c(x_i) \in H^2(X_i[s], M_X)$ associated to the line bundle $\mathcal{O}(x_i)$ (cf. [Mochizuki], Proposition 1.6, and Proposition 1.8 (iii)). Thus, the pro-Chern class $\hat{c}(s)$ is a Picard element. More precisely, it is the “pro-Picard element” induced by the $(\mathcal{O}(x_i))_{i \geq 1}$.

If the above condition is satisfied one says that the Chern class of the section s is algebraic, or that the section s is well-behaved (cf. [Saïdi], Proposition 1.5.1).

One can easily prove that if the section s is well-behaved, then the section s is good in the sense of Definition 3.6 (cf. [Saïdi], Proposition 1.5.2).

§4. The Cuspidalisation of Sections of Arithmetic Fundamental Groups.

In this section we introduce the problem of cuspidalisation of group-theoretic sections of arithmetic fundamental groups.

We follow the notations in §1.

In particular, $\Sigma \subseteq \mathfrak{Primes}$ is a non-empty set of prime integers, and we have the natural exact sequence

$$(1.2) \quad 1 \rightarrow \Delta_X \rightarrow \Pi_X \xrightarrow{\text{pr}_X} G_k \rightarrow 1,$$

where Π_X is the geometrically pro- Σ arithmetic fundamental group of X .

4.1. In this sub-section we recall the definition of (geometrically) cuspidally central, and cuspidally abelian, arithmetic fundamental groups of affine hyperbolic curves, and the definition of cupidally abelian absolute Galois groups of function fields of curves.

4.1.1. Let $U \subseteq X$ be a non-empty open subscheme of X . The geometric point η of X determines naturally a geometric point η of U , and a geometric point $\bar{\eta}$ of $\bar{U} \stackrel{\text{def}}{=} U \times_k \bar{k}$. Write

$$\Delta_U \stackrel{\text{def}}{=} \pi_1(\bar{U}, \bar{\eta})^\Sigma$$

for the maximal pro- Σ quotient of the fundamental group $\pi_1(\bar{U}, \bar{\eta})$ of \bar{U} with base point η , and

$$\Pi_U \stackrel{\text{def}}{=} \pi_1(U, \eta) / \text{Ker}(\pi_1(\bar{U}, \bar{\eta}) \twoheadrightarrow \pi_1(\bar{U}, \bar{\eta})^\Sigma)$$

for the quotient of the arithmetic fundamental group $\pi_1(U, \eta)$ by the kernel of the natural surjective homomorphism $\pi_1(\bar{U}, \bar{\eta}) \twoheadrightarrow \pi_1(\bar{U}, \bar{\eta})^\Sigma$, which is a normal subgroup of $\pi_1(U, \eta)$.

Thus, we have a natural exact sequence

$$1 \rightarrow \Delta_U \rightarrow \Pi_U \xrightarrow{\text{pr}_{U, \Sigma}} G_k \rightarrow 1,$$

which fits into the following commutative diagram:

$$\begin{array}{ccccccc} 1 & \longrightarrow & \Delta_U & \longrightarrow & \Pi_U & \xrightarrow{\text{pr}_{U, \Sigma}} & G_k \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \text{id} \downarrow \\ 1 & \longrightarrow & \Delta_X & \longrightarrow & \Pi_X & \xrightarrow{\text{pr}_{X, \Sigma}} & G_k \longrightarrow 1 \end{array}$$

where the left, and middle, vertical homomorphisms are surjective, and are naturally induced by the natural surjective homomorphisms $\pi_1(\bar{U}, \bar{\eta}) \twoheadrightarrow \pi_1(\bar{X}, \bar{\eta})$, and $\pi_1(U, \eta) \twoheadrightarrow \pi_1(X, \eta)$.

Let

$$I_U \stackrel{\text{def}}{=} \text{Ker}(\Pi_U \twoheadrightarrow \Pi_X) = \text{Ker}(\Delta_U \twoheadrightarrow \Delta_X).$$

We shall refer to I_U as the cuspidal subgroup of Π_U (cf. [Mochizuki], Definition 1.5). It is the normal subgroup of Π_U generated by the (pro- Σ) inertia subgroups at the geometric points of $S \stackrel{\text{def}}{=} X \setminus U$. We have the following natural exact sequence

$$(4.1) \quad 1 \rightarrow I_U \rightarrow \Pi_U \rightarrow \Pi_X \rightarrow 1.$$

Let I_U^{ab} be the maximal abelian quotient of I_U . By pushing out the exact sequence (4.1) by the natural surjective homomorphism $I_U \twoheadrightarrow I_U^{\text{ab}}$, we obtain a natural commutative diagram:

$$\begin{array}{ccccccc} 1 & \longrightarrow & I_U & \longrightarrow & \Pi_U & \longrightarrow & \Pi_X \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \text{id} \downarrow \\ 1 & \longrightarrow & I_U^{\text{ab}} & \longrightarrow & \Pi_U^{\text{c-ab}} & \longrightarrow & \Pi_X \longrightarrow 1 \end{array}$$

We will refer to the quotient $\Pi_U^{c-\text{ab}}$ of Π_U as the maximal cuspidally abelian quotient of Π_U , with respect to the natural homomorphism $\Pi_U \twoheadrightarrow \Pi_X$ (cf. [Mochizuki], Definition 1.5).

Similarly, we can define the maximal cuspidally abelian quotient $\Delta_U^{c-\text{ab}}$ of Δ_U , with respect to the natural homomorphism $\Delta_U \twoheadrightarrow \Delta_X$, which sits in a natural exact sequence

$$(4.2) \quad 1 \rightarrow I_U^{\text{ab}} \rightarrow \Delta_U^{c-\text{ab}} \rightarrow \Delta_X \rightarrow 1.$$

Write I_U^{cn} for the maximal quotient of I_U^{ab} on which the action of Δ_X , which is naturally deduced from the exact sequence (4.2), is trivial. By pushing out the sequence (4.2) by the natural surjective homomorphism $I_U^{\text{ab}} \twoheadrightarrow I_U^{\text{cn}}$, we obtain a natural exact sequence

$$(4.3) \quad 1 \rightarrow I_U^{\text{cn}} \rightarrow \Delta_U^{c-\text{cn}} \rightarrow \Delta_X \rightarrow 1.$$

Define

$$\Pi_U^{c-\text{cn}} \stackrel{\text{def}}{=} \Pi_U^{c-\text{ab}} / \text{Ker}(I_U^{\text{ab}} \twoheadrightarrow I_U^{\text{cn}}),$$

which sits naturally in the following exact sequence

$$(4.4) \quad 1 \rightarrow I_U^{\text{cn}} \rightarrow \Pi_U^{c-\text{cn}} \rightarrow \Pi_X \rightarrow 1.$$

We shall refer to the quotient $\Pi_U^{c-\text{cn}}$ of Π_U as the maximal (geometrically) cuspidally central quotient of Π_U , with respect to the natural homomorphism $\Pi_U \twoheadrightarrow \Pi_X$ (cf. loc. cit.).

We have a natural commutative diagram of exact sequences:

$$\begin{array}{ccccccccc} 1 & \longrightarrow & I_U & \longrightarrow & \Pi_U & \longrightarrow & \Pi_X & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \text{id} \downarrow & & \\ 1 & \longrightarrow & I_U^{\text{ab}} & \longrightarrow & \Pi_U^{c-\text{ab}} & \longrightarrow & \Pi_X & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \text{id} \downarrow & & \\ 1 & \longrightarrow & I_U^{\text{cn}} & \longrightarrow & \Pi_U^{c-\text{cn}} & \longrightarrow & \Pi_X & \longrightarrow & 1 \end{array}$$

4.1.2. Similarly, we have a natural exact sequence of absolute Galois groups

$$1 \rightarrow G_{\bar{k}.K_X} \rightarrow G_{K_X} \rightarrow G_k \rightarrow 1,$$

where $G_{\bar{k}.K_X} \stackrel{\text{def}}{=} \text{Gal}(K_X^{\text{sep}}/\bar{k}.K_X)$, and $G_{K_X} \stackrel{\text{def}}{=} \text{Gal}(K_X^{\text{sep}}/K_X)$.

Let

$$\overline{G}_X \stackrel{\text{def}}{=} G_{\bar{k}.K_X}^{\Sigma}$$

be the maximal pro- Σ quotient of $G_{\bar{k}.K_X}$, and

$$G_X \stackrel{\text{def}}{=} G_{K_X} / \text{Ker}(G_{\bar{k}.K_X} \twoheadrightarrow \overline{G}_X),$$

which insert into the following commutative diagram of exact sequences:

$$\begin{array}{ccccccc}
1 & \longrightarrow & \overline{G}_X & \longrightarrow & G_X & \xrightarrow{\tilde{\text{pr}}_{X,\Sigma}} & G_k \longrightarrow 1 \\
& & \downarrow & & \downarrow & & \text{id} \downarrow \\
1 & \longrightarrow & \Delta_X & \longrightarrow & \Pi_X & \xrightarrow{\text{pr}_{X,\Sigma}} & G_k \longrightarrow 1
\end{array}$$

where the left vertical maps are the natural surjective homomorphisms.

Let

$$I_X \stackrel{\text{def}}{=} \text{Ker}(G_X \twoheadrightarrow \Pi_X) = \text{Ker}(\overline{G}_X \twoheadrightarrow \Delta_X).$$

We will refer to I_X as the cuspidal subgroup of G_X . It is the normal subgroup of G_X generated by the (pro- Σ) inertia subgroups at all geometric closed points of X . We have the following natural exact sequence

$$1 \rightarrow I_X \rightarrow G_X \rightarrow \Pi_X \rightarrow 1.$$

Let I_X^{ab} be the maximal abelian quotient of I_X . By pushing out the above sequence by the natural surjective homomorphism $I \twoheadrightarrow I^{\text{ab}}$, we obtain a natural exact sequence

$$(4.5) \quad 1 \rightarrow I^{\text{ab}} \rightarrow G_X^{\text{c-ab}} \rightarrow \Pi_X \rightarrow 1.$$

We will refer to the quotient $G_X^{\text{c-ab}}$ as the maximal cuspidally abelian quotient of G_X , with respect to the natural homomorphism $G_X \twoheadrightarrow \Pi_X$. Note that $G_X^{\text{c-ab}}$ is naturally identified with the projective limit

$$\varprojlim_U \Pi_U^{\text{c-ab}},$$

where the limit runs over all open subschemes U of X .

4.2. Next, we consider a continuous group-theoretic section $s : G_k \rightarrow \Pi_X$ of the natural projection $\text{pr}_X : \Pi_X \twoheadrightarrow G_k$.

Definition 4.3 (Lifting of Group-Theoretic Sections). Let $U \subseteq X$ be a non-empty open subscheme. We say that a continuous group-theoretic section $s_U : G_k \rightarrow \Pi_U$, of the natural projection $\text{pr}_U \stackrel{\text{def}}{=} \text{pr}_{U,\Sigma} : \Pi_U \twoheadrightarrow G_k$ (meaning that $\text{pr}_U \circ s_U = \text{id}_{G_k}$), is a lifting of the section $s : G_k \rightarrow \Pi_X$, if s_U fits into a commutative diagram:

$$\begin{array}{ccc}
G_k & \xrightarrow{s_U} & \Pi_U \\
\text{id} \downarrow & & \downarrow \\
G_k & \xrightarrow{s} & \Pi_X
\end{array}$$

where the right vertical homomorphism is the natural one.

More generally, we say that a group-theoretic section $\tilde{s} : G_k \rightarrow G_X$ of the natural projection $\tilde{\text{pr}} \stackrel{\text{def}}{=} \tilde{\text{pr}}_{X,\Sigma} : G_X \twoheadrightarrow G_k$ (meaning that $\tilde{\text{pr}} \circ \tilde{s} = \text{id}_{G_k}$) is a lifting of the section s , if \tilde{s} fits into a commutative diagram:

$$\begin{array}{ccc}
G_k & \xrightarrow{\tilde{s}} & G_X \\
\text{id} \downarrow & & \downarrow \\
G_k & \xrightarrow{s} & \Pi_X
\end{array}$$

In connection with the Grothendieck anabelian section conjecture, it is natural to consider the following problem.

The Cuspidalisation Problem for sections of Arithmetic Fundamental Groups. Given a group-theoretic section $s : G_k \rightarrow \Pi_X$ as above, and a non-empty open subscheme $U \subseteq X$, is it possible to construct a lifting $s_U : G_k \rightarrow \Pi_U$ of s ? Similarly, given a group-theoretic section $s : G_k \rightarrow \Pi_X$ as above, is it possible to construct a lifting $\tilde{s} : G_k \rightarrow G_X$ of s ?

Remarks 4.4.

(i) One can easily verify that if the section s is point-theoretic, then the section s can be lifted to a section $s_U : G_k \rightarrow \Pi_U$ of the natural projection $\Pi_U \twoheadrightarrow G_k$, for every open subscheme $U \subseteq X$, and can also be lifted to a section $\tilde{s} : G_k \rightarrow G_X$ of the natural projection $G_X \twoheadrightarrow G_k$.

(ii) Note that a positive solution to the cuspidalisation problem, in the case where k is finitely generated over \mathbb{Q} (resp. p -adic local field), and $\Sigma = \mathfrak{Primes}$, plus a positive solution to the BGASC (resp. p -adic BGASC), gives a positive solution to the GASC (resp. p -adic GASC), as follows easily from Lemma 2.5.

Our main result concerning the cuspidalisation problem is the following, which shows that good sections of arithmetic fundamental groups behave well with respect to this problem.

Before stating our result, we recall some definitions.

Definition 4.5. (i) We say that the field k is slim if its absolute Galois group G_k is slim in the sense of [Mochizuki], §0, meaning that every open subgroup of G_k is centre free. Examples of slim fields include number fields, and p -adic local fields (cf. [Mochizuki2], Theorem 1.1.1).

One defines in a similar way the notion of a slim profinite group G , meaning that every open subgroup of G is centre free.

(ii) We say that the field k is Σ -regular, if for every prime integer $l \in \Sigma$, and every finite extension k'/k , the l -part of the cyclotomic character $\chi_l : G_{k'} \rightarrow \mathbb{Z}_l^\times$ is not trivial; or equivalently, if for every prime integer $l \in \Sigma$ the image of the l -part of the cyclotomic character $\chi_l : G_k \rightarrow \mathbb{Z}_l^\times$ is infinite.

Examples of Σ -regular fields (for every non-empty set Σ of prime integers) include number fields, p -adic local fields, and finite fields.

The field k is Σ -regular if and only if, for every finite extension k'/k , the $G_{k'}$ -module M_X has no non trivial fixed elements.

The following is our main result on the cuspidalisation problem (cf. [Saïdi], Theorem 2.6).

Theorem 4.6 (Lifting of Uniformly Good Sections to Cuspidally abelian Arithmetic Fundamental Groups over Slim Fields). *Assume that the section $s : G_k \rightarrow \Pi_X$ is uniformly good (in the sense of Definition 3.6), and that the field k is slim, and Σ -regular (cf. Definition 4.5). Let $U \subseteq X$ be a non-empty open subscheme of X , and $\Pi_U^{\text{c-ab}}$ the maximal cuspidally abelian quotient of Π_U , with respect to the natural surjective homomorphism $\Pi_U \twoheadrightarrow \Pi_X$.*

Then there exists a section $s_U^{\text{c-ab}} : G_k \rightarrow \Pi_U^{\text{c-ab}}$ of the natural projection $\Pi_U^{\text{c-ab}} \twoheadrightarrow G_k$ which lifts the section s , i.e. which inserts into the following commu-

tative diagram:

$$\begin{array}{ccc} G_k & \xrightarrow{s_U^{c-ab}} & \Pi_U^{c-ab} \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s} & \Pi_X \end{array}$$

Moreover, one can construct for every non-empty open subscheme $U \stackrel{\text{def}}{=} X \setminus S$ of X a section $s_U^{c-ab} : G_k \rightarrow \Pi_U^{c-ab}$ as above (i.e. which lifts the section s), such that for every non-empty open subscheme $V \stackrel{\text{def}}{=} X \setminus T$ of X , with $U \subseteq V$, we have the following commutative diagram:

$$\begin{array}{ccc} G_k & \xrightarrow{s_U^{c-cn}} & \Pi_U^{c-ab} \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s_V^{c-cn}} & \Pi_V^{c-ab} \end{array}$$

where the right vertical homomorphism is the natural one.

As a corollary of the above result one obtains the following.

Corollary 4.7 (Lifting of Uniformly Good Sections to Cuspidally abelian Galois Groups over Slim Fields). *Assume that the field k is slim, and Σ -regular (cf. Definition 4.5). Let $s : G_k \rightarrow \Pi_X$ be a uniformly good group-theoretic section of the natural projection $\Pi_X \twoheadrightarrow G_k$ (in the sense of Definition 3.6). Then there exists a section $s^{c-ab} : G_k \rightarrow G_X^{c-ab}$ of the natural projection $G_X^{c-ab} \twoheadrightarrow G_k$, which lifts the section s , i.e. which inserts into the following commutative diagram:*

$$\begin{array}{ccc} G_k & \xrightarrow{s^{c-ab}} & G_X^{c-ab} \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s} & \Pi_X \end{array}$$

4.8. Next, we would like to explain the basic idea behind the proof of Theorem 4.6. See loc. cit. for more details.

Assume that the field k is slim, Σ -regular, and the section $s : G_k \rightarrow \Pi_X$ is uniformly good (in the sense of Definition 3.6).

Let $U \stackrel{\text{def}}{=} X \setminus S$ be a non-empty open subscheme of X , and Π_U^{c-cn} the maximal (geometrically) cuspidally central quotient of Π_U , with respect to the natural homomorphism $\Pi_U \twoheadrightarrow \Pi_X$.

One would like to show, in a first step, that there exists a section $s_U^{c-cn} : G_k \rightarrow \Pi_U^{c-cn}$ of the natural projection $\Pi_U^{c-cn} \twoheadrightarrow G_k$ which lifts the section s , i.e. which inserts into the following commutative diagram:

$$\begin{array}{ccc} G_k & \xrightarrow{s_U^{c-cn}} & \Pi_U^{c-cn} \\ \text{id} \downarrow & & \downarrow \\ G_k & \xrightarrow{s} & \Pi_X \end{array}$$

First, one treats the case where the set $S = \{x_i\}_{i=1}^n \subseteq X(k)$ consists of finitely many k -rational points. We will assume, without loss of generality, that $S = \{x\}$ consists of a single rational point $x \in X(k)$.

The maximal (geometrically) cuspidally central quotient $\Pi_{U_x}^{c-\text{cn}}$ of Π_{U_x} , with respect to the natural projection $\Pi_{U_x} \rightarrow \Pi_X$, sits naturally in the following exact sequence

$$(4.6) \quad 1 \rightarrow M_X \rightarrow \Pi_{U_x}^{c-\text{cn}} \rightarrow \Pi_X \rightarrow 1$$

(cf. [Mochizuki], Proposition 1.8).

By pulling back the group extension (4.6) by the section $s : G_k \rightarrow \Pi_X$, we obtain a group extension

$$(4.7) \quad 1 \rightarrow M_X \rightarrow s^*(\Pi_{U_x}^{c-\text{cn}}) \rightarrow G_k \rightarrow 1,$$

which inserts naturally in the following commutative diagram:

$$\begin{array}{ccccccc} 1 & \longrightarrow & M_X & \longrightarrow & s^*(\Pi_{U_x}^{c-\text{cn}}) & \longrightarrow & G_k \longrightarrow 1 \\ & & \text{id} \downarrow & & \downarrow & & s \downarrow \\ 1 & \longrightarrow & M_X & \longrightarrow & \Pi_{U_x}^{c-\text{cn}} & \longrightarrow & \Pi_X \longrightarrow 1 \end{array}$$

where the right square is cartesian.

The class in $H^2(\Pi_X, M_X)$ of the group extension (4.6) coincides, via the natural identification $H^2(\Pi_X, M_X) \xrightarrow{\sim} H^2(X, M_X)$, with the étale Chern class $c(x) \in H^2(\Pi_X, M_X)$ associated to the degree 1 line bundle $\mathcal{O}(x)$ (cf. [Mochizuki3], lemma 4.2). The class in $H^2(G_k, M_X)$ of the group extension (4.7) coincides then with the image $s^*(c(x))$ of the Chern class $c(x)$ via the (restriction) homomorphism $s^* : H^2(X, M_X) \rightarrow H^2(G_k, M_X)$, which is naturally induced by s .

This image equals 0, since the section s is assumed to be good. This follows from the very definition of goodness (cf. Definition 3.6). Thus, the group extension (4.7) admits group-theoretic splittings. A splitting of the group extension (4.7) determines a section $s_U^{c-\text{cn}} : G_k \rightarrow \Pi_U^{c-\text{cn}}$, which lifts the section s .

The case where the set S consists of finitely many, not necessarily rational, points is treated in a similar way by using a descent argument, which resorts to the slimness of k , and the fact that k is Σ -regular.

In general, and in order to lift the section s to a section $s_U^{c-\text{ab}} : G_k \rightarrow \Pi_U^{c-\text{ab}}$, one uses the following description of $\Pi_U^{c-\text{ab}}$.

For a finite étale Galois cover $X' \rightarrow X$, with Galois group $\text{Gal}(X'/X)$, let $U' \stackrel{\text{def}}{=} U \times_X X'$, and $\Pi_{U'}^{c-\text{cn}}$ the maximal (geometrically) cuspidally central quotient of $\Pi_{U'}$, with respect to the natural surjective homomorphism $\Pi_{U'} \rightarrow \Pi_{X'}$, which is slim.

Denote by $\Pi_{U'}^{c-\text{cn}} \rtimes^{\text{out}} \text{Gal}(X'/X)$ the profinite group which is obtained by pulling back the exact sequence

$$1 \rightarrow \Pi_{U'}^{c-\text{cn}} \rightarrow \text{Aut}(\Pi_{U'}^{c-\text{cn}}) \rightarrow \text{Out}(\Pi_{U'}^{c-\text{cn}}) \rightarrow 1,$$

by the natural homomorphism:

$$\text{Gal}(X'/X) \rightarrow \text{Out}(\Pi_{U'}^{c-\text{cn}}).$$

Thus, we have a natural exact sequence:

$$1 \rightarrow \Pi_{U'}^{\text{c-cn}} \rightarrow \Pi_{U'}^{\text{c-cn}} \rtimes^{\text{out}} \text{Gal}(X'/X) \rightarrow \text{Gal}(X'/X) \rightarrow 1.$$

which inserts into the following commutative diagram:

$$\begin{array}{ccccccc}
& 1 & & 1 & & & \\
& \downarrow & & \downarrow & & & \\
& I_{U'}^{\text{c-cn}} & \xrightarrow{\text{id}} & I_{U'}^{\text{c-cn}} & & & \\
& \downarrow & & \downarrow & & & \\
1 & \longrightarrow & \Pi_{U'}^{\text{c-cn}} & \longrightarrow & \Pi_{U'}^{\text{c-cn}} \rtimes^{\text{out}} \text{Gal}(X'/X) & \longrightarrow & \text{Gal}(X'/X) \longrightarrow 1 \\
& \downarrow & & \downarrow & & \text{id} \downarrow & \\
1 & \longrightarrow & \Pi_{X'} & \longrightarrow & \Pi_X & \longrightarrow & \text{Gal}(X'/X) \longrightarrow 1 \\
& \downarrow & & \downarrow & & \downarrow & \\
& 1 & & 1 & & 1 &
\end{array}$$

Then we have a natural isomorphism:

$$\Pi_U^{\text{c-ab}} \xrightarrow{\sim} \varprojlim_{X' \rightarrow X} \Pi_{U'}^{\text{c-cn}} \rtimes^{\text{out}} \text{Gal}(X'/X),$$

where the projective limit is taken over all finite étale Galois cover $X' \rightarrow X$ (cf. [Saïdi], Proposition 2.5).

§5. Applications to the Grothendieck Anabelian Section Conjecture.

In this section we state our main applications of the results concerning the cuspidalisation problem for sections of arithmetic fundamental groups, in §4, to the Grothendieck anabelian section conjecture.

As we already mentioned (cf. Remarks 4.4, (ii)), a positive answer to the cuspidalisation problem, in the case where k is finitely generated over \mathbb{Q} (resp. p -adic local field), and $\Sigma = \mathfrak{Primes}$, plus a positive answer to the BGASC (resp. p -adic BGASC), implies a positive answer to the GASC (resp. p -adic GASC).

The following result of Koenigsmann concerning the p -adic BGASC is fundamental (cf. [Koenigsmann]).

Theorem 5.1 (Koenigsmann). *The p -adic BGASC holds true. More precisely, assume that k is a p -adic local field, and $\Sigma = \mathfrak{Primes}$. Let $s : G_k \rightarrow G_X$ be a group-theoretic section of the natural projection $G_X \twoheadrightarrow G_k$. Then the image $s(G_k)$ is contained in a decomposition subgroup D_x associated to a unique rational point $x \in X(k)$. In particular, $X(k) \neq \emptyset$.*

This result has been strengthened by Pop, who proved the following (see [Pop]). For a profinite group H , and a prime integer p , we denote by H'' the maximal $\mathbb{Z}/p\mathbb{Z}$ -metabelian quotient of H . Thus, H'' is the second quotient of the $\mathbb{Z}/p\mathbb{Z}$ -derived series of H .

Theorem 5.2. (Pop). *Assume that k is a p -adic local field, which contains a primitive p -th root of 1, and assume $p \in \Sigma$. Let $s : G''_k \rightarrow G''_{K_X}$ be a group-theoretic section of the natural projection $G''_{K_X} \twoheadrightarrow G''_k$. Then the image $s(G''_k)$ is contained in a decomposition subgroup $D_x \subset G''_{K_X}$ associated to a unique rational point $x \in X(k)$. In particular, $X(k) \neq \emptyset$. Here the $(\)''$ of the various profinite groups are with respect to the prime p , i.e. the second quotients of the $\mathbb{Z}/p\mathbb{Z}$ -derived series.*

The above Theorem of Pop can be viewed as a very “minimalistic” version of the birational Grothendieck anabelian section conjecture over p -adic local fields. Note that the quotient G''_k of G_k is finite in this case.

Let us mention few words on the proof of the above result of Pop.

The technical tool in the proof, which produces a rational point $x \in X(k)$ starting from a section $s : G''_k \rightarrow G''_{K_X}$, is the following.

For a profinite group H , denote by H' the maximal quotient of H which is abelian and annihilated by p . Thus, H' is the first quotient of the $\mathbb{Z}/p\mathbb{Z}$ -derived series of H .

The existence of the section $s : G''_k \rightarrow G''_{K_X}$ implies the existence of a section $s' : G'_k \rightarrow G'_{K_X}$ of the natural projection $G'_{K_X} \twoheadrightarrow G'_k$.

Let \tilde{L}/K_X be the subextension of K_X^{sep}/K_X with Galois group G'_{K_X} , and L/K_X the subextension of \tilde{L}/K_X corresponding to the subgroup $s'(G'_k)$ of G'_{K_X} . Then L is a field of transcendence degree 1 over k . For such a field, Pop proved a local-global principle for Brauer groups, which generalises a similar principle for function fields of curves over p -adic local fields due to Lichtenbaum, and which reads as follows.

The natural diagonal homomorphism

$$\text{Br}(L) \rightarrow \prod_v \text{Br}(L_v),$$

where the product runs over all rank 1 valuations v of L , is injective.

Now the existence of the section $s : G''_k \rightarrow G''_{K_X}$ implies that the natural homomorphism

$$\text{Br}(k) \rightarrow \text{Br}(L)$$

is injective (see [Pop] for more details).

Let $\alpha \in \text{Br}(k)$ be an element of order p . Then α survives in $\text{Br}(L_v)$, i.e. its image is non-zero, for some rank 1 valuation v of L , by the above local-global principle. Pop then proves (the proof is rather technical) that v is the valuation associated to a unique k -rational point $x \in X(k)$.

5.3. Using the technique of cuspidalisation of sections of arithmetic fundamental groups, introduced in §4, one can hope to prove the p -adic GASC by reducing it to the p -adic version of the BGASC which was proved by Pop. We are able to prove that this is indeed the case, under some additional assumptions.

Before stating our result we will define the notion of good sections of cuspidally abelian absolute Galois groups.

Assume that k is a p -adic local field, and $p \in \Sigma$. Let G_X be the geometrically pro- Σ absolute Galois group of K_X , and $G_X^{\text{c-ab}}$ the maximal cuspidally abelian quotient of G_X , with respect to the natural surjective homomorphism $G_X \twoheadrightarrow \Pi_X$ (cf. 4.1.2).

Let

$$\tilde{s} : G_k \rightarrow G_X^{c-\text{ab}}$$

be a continuous group-theoretic section of the natural projection $G_X^{c-\text{ab}} \twoheadrightarrow G_k$. Let \tilde{L}/K_X be the subextension of K_X^{sep}/K_X with Galois group $G_X^{c-\text{ab}}$, and L/K_X the subextension of \tilde{L}/K_X corresponding to the closed subgroup $\tilde{s}(G_k)$ of $G_X^{c-\text{ab}}$.

We say that the section \tilde{s} is a good, or tame point-theoretic, group-theoretic section, if the natural homomorphism

$$\text{Br}(k) \rightarrow \text{Br}(L)$$

is injective (cf. [Saïdi], 1.7, for more details).

Our main result concerning the Grothendieck anabelian section conjecture over p -adic local fields is the following.

Theorem 5.4. *Assume that k is a p -adic local field, and $p \in \Sigma$. Let $s : G_k \rightarrow \Pi_X$ be a group-theoretic section of the natural projection $\Pi_X \twoheadrightarrow G_k$. Assume that s is a good section (in the sense of Definition 3.6). Then there exists a section $s^{c-\text{ab}} : G_k \rightarrow G_X^{c-\text{ab}}$ of the natural projection $G_X^{c-\text{ab}} \twoheadrightarrow G_k$, which lifts the section s . Furthermore, if the section $s^{c-\text{ab}} : G_k \rightarrow G_X^{c-\text{ab}}$ is a good section (in the sense of 5.3), then $X(k) \neq \emptyset$.*

Proof. The first assertion is Corollary 4.7.

For a profinite group H , and a prime integer p , denote by H' the maximal quotient of H which is abelian, and annihilated by p . Thus, H' is the first quotient of the $\mathbb{Z}/p\mathbb{Z}$ -derived series of H .

The existence of the section $s^{c-\text{ab}} : G_k \rightarrow G_X^{c-\text{ab}}$ implies the existence of a section $s' : G'_k \rightarrow G'_X$ of the natural projection $G'_X \twoheadrightarrow G'_k$.

Let \tilde{L}/K_X be the sub-extension of K_X^{sep}/K_X which corresponds to the closed subgroup $\text{Ker}(G_X \twoheadrightarrow G'_X)$ of G_X , and L/K_X the sub-extension of \tilde{L}/K_X which corresponds to the closed subgroup $s'(G'_k)$ of G'_X .

Assume that the section $s^{c-\text{ab}} : G_k \rightarrow G_X^{c-\text{ab}}$ is good (in the sense of 5.3). Then the natural homomorphism $\text{Br } k \rightarrow \text{Br } L$ is injective. Under this assumption (which is implied, in the framework of the proof by Pop of Theorem 5.2, by the lifting property of the section $s' : G'_k \rightarrow G'_X$ to a section $s'' : G''_k \rightarrow G''_X$ which is imposed in [Pop]) Pop proves that the image $s'(G'_k)$ is contained in a decomposition subgroup $D_x \subset G'_{K_X}$ associated to a unique rational point $x \in X(k)$ (cf. loc. cit). In particular, $X(k) \neq \emptyset$ in this case. \square

With the same notations as in theorem 5.4, the author expects that if s is a good section, then any lifting $s^{c-\text{ab}} : G_k \rightarrow G_X^{c-\text{ab}}$ of s (which exists by Corollary 4.7) is automatically good in the sense of 5.3. In particular, following Theorem 5.4, every good section s should be point-theoretic, if $p \in \Sigma$. The author is unable to prove this at the moment of writing this paper.

5.5. One can prove, using cuspidalisation techniques of sections of arithmetic fundamental groups, the following (unconditional) version of the p -adic GASC.

Theorem 5.6. *Assume that k is a p -adic local field, and $\Sigma = \{p\}$. Let $S \subset X$ be a set of closed points of X which is uniformly dense in X for the p -adic topology,*

meaning that for each finite extension k'/k , $S(k')$ is dense in $X(k')$ for the p -adic topology. Write $\Pi_{X \setminus S}$ for the geometrically pro- Σ quotient of the arithmetic fundamental group $\pi_1(X \setminus S, \eta)$ (which is defined in a similar way as Π_X , cf. 1.1).

Let $s : G_k \rightarrow \Pi_{X \setminus S}$ be a continuous group-theoretic section of the natural projection $\Pi_{X \setminus S} \twoheadrightarrow G_k$. Then s is point-theoretic, i.e. the image $s(G_k)$ in $\Pi_{X \setminus S}$ is a decomposition group $D_x \subset \Pi_{X \setminus S}$, associated to unique rational point $x \in X(k)$.

An example of a set S satisfying the assumptions of Theorem 5.6 is the set of algebraic points, in the case where X is defined over a number field

A proof of Theorem 5.6, was communicated orally to the author by A. Tamagawa. The proof relies on the idea of cuspidalisation, and consists in showing that the section s can be lifted to a section $\tilde{s} : G_k \rightarrow G_X$ of the natural projection $G_X \twoheadrightarrow G_k$, where G_X is the geometrically pro- Σ absolute Galois group of K_X . One then reduces the proof to the p -adic version of the BGASC that was proven by Pop.

The proof uses the fact that the Galois group G_k of k is topologically finitely generated, and consists in showing that given a (not necessarily geometrically connected) ramified Galois cover $Y \rightarrow X$ with Galois group G , one can "approximate" it by a Galois cover $Y' \rightarrow X$ with Galois group G , which is ramified only above points which are contained in S . The later relies on an approximation argument, à la Artin, on Hurwitz spaces of covers.

5.7. In [Esnault-Wittenberg1] sections of geometrically abelian absolute Galois groups of function fields of curves over number fields were investigated.

It is shown in loc. cit. that the existence of such sections implies (in fact is equivalent to) the existence of degree 1 divisors on the curve, under a finiteness condition of the Tate-Shafarevich group of the jacobian of the curve.

As an application of our results on the cuspidalisation of sections of arithmetic fundamental groups, we can prove an analogous result for good sections of arithmetic fundamental groups.

Our main result concerning the Grothendieck anabelian section conjecture over number fields is the following.

Theorem 5.8. *Assume that k is a number field, and $\Sigma = \mathfrak{P}\text{rimes}$. Let $s : G_k \rightarrow \Pi_X$ be a group-theoretic section of the natural projection $\Pi_X \twoheadrightarrow G_k$. Assume that s is a uniformly good section in the sense of definition 3.6, and that the jacobian variety of X has a finite Tate-Shafarevich group. Then there exists a divisor of degree 1 on X .*

Proof. Follows formally from Corollary 4.7, and Theorem 2.1 in [Esnault-Wittenberg1]. \square

§6. On a Weak Form of the p -adic Grothendieck Anabelian section Conjecture.

In this section we discuss a weak form of the p -adic GASC. We will use the following notations.

Let $p > 0$ be a fixed prime integer. Let k be a p -adic local field, i.e. k is a finite extension of \mathbb{Q}_p , \mathcal{O}_k its ring of integers, and F its residue field.

Let X be a proper, smooth, geometrically connected, and hyperbolic curve over k .

For a non-empty set of prime integers $\Sigma \subseteq \mathfrak{P}\text{rimes}$, write Π_X for the geometrically pro- Σ fundamental group of X , which sits in the exact sequence

$$1 \rightarrow \Delta_X \rightarrow \Pi_X \rightarrow G_k \rightarrow 1,$$

where Δ_X is the maximal pro- Σ quotient of the fundamental group of $\overline{X} \stackrel{\text{def}}{=} X \times_k \bar{k}$ (cf. 1.1).

Recall the natural map (cf. 2.1)

$$\varphi_X \stackrel{\text{def}}{=} \varphi_{X,\Sigma} : X(k) \rightarrow \overline{\text{Sec}}_{\Pi_X}.$$

6.1. Assume that the hyperbolic k -curve X has good reduction over O_k , i.e. X extends to a smooth, proper, and relative curve \mathcal{X} over O_k , and $p \notin \Sigma$. Let $\mathcal{X}_s \stackrel{\text{def}}{=} \mathcal{X} \times_{O_k} F$ be the special fibre of \mathcal{X} . Let ξ be a geometric point of \mathcal{X}_s above the generic point of \mathcal{X}_s . Then ξ determines naturally an algebraic closure \overline{F} of F , and a geometric point $\bar{\xi}$ of $\overline{\mathcal{X}}_s \stackrel{\text{def}}{=} \mathcal{X}_s \times_F \overline{F}$.

There exists a natural exact sequence of profinite groups

$$1 \rightarrow \pi_1(\overline{\mathcal{X}}_s, \bar{\xi}) \rightarrow \pi_1(\mathcal{X}_s, \xi) \xrightarrow{\text{pr}} G_F \rightarrow 1.$$

Here $\pi_1(\mathcal{X}_s, \xi)$ denotes the arithmetic étale fundamental group of \mathcal{X}_s with base point ξ , $\pi_1(\overline{\mathcal{X}}_s, \bar{\xi})$ the étale fundamental group of $\overline{\mathcal{X}}_s \stackrel{\text{def}}{=} \mathcal{X}_s \times_F \overline{F}$ with base point $\bar{\xi}$, and $G_F \stackrel{\text{def}}{=} \text{Gal}(\overline{F}/F)$ the absolute Galois group of F .

Write

$$\Delta_{\mathcal{X}_s} \stackrel{\text{def}}{=} \pi_1(\overline{\mathcal{X}}_s, \bar{\xi})^\Sigma$$

for the maximal pro- Σ quotient of $\pi_1(\overline{\mathcal{X}}_s, \bar{\xi})$, and

$$\Pi_{\mathcal{X}_s} \stackrel{\text{def}}{=} \pi_1(\mathcal{X}_s, \xi) / \text{Ker}(\pi_1(\overline{\mathcal{X}}_s, \bar{\xi}) \twoheadrightarrow \pi_1(\overline{\mathcal{X}}_s, \bar{\xi})^\Sigma)$$

for the quotient of $\pi_1(\mathcal{X}_s, \xi)$ by the kernel of the natural surjective homomorphism $\pi_1(\overline{\mathcal{X}}_s, \bar{\xi}) \twoheadrightarrow \pi_1(\overline{\mathcal{X}}_s, \bar{\xi})^\Sigma$, which is a normal subgroup of $\pi_1(\mathcal{X}_s, \xi)$.

Thus, we have an exact sequence of profinite groups

$$(6.1) \quad 1 \rightarrow \Delta_{\mathcal{X}_s} \rightarrow \Pi_{\mathcal{X}_s} \xrightarrow{\text{pr}} G_F \rightarrow 1.$$

Moreover, after a suitable choice of the base points ξ , and η , there exists a natural commutative specialisation diagram:

$$(6.2) \quad \begin{array}{ccccccc} 1 & \longrightarrow & \Delta_X & \longrightarrow & \Pi_X & \longrightarrow & G_k \longrightarrow 1 \\ & & \downarrow & & \text{Sp}_X \downarrow & & \downarrow \\ 1 & \longrightarrow & \Delta_{\mathcal{X}_s} & \longrightarrow & \Pi_{\mathcal{X}_s} & \longrightarrow & G_F \longrightarrow 1 \end{array}$$

where the left vertical homomorphism $\text{Sp} : \Delta_X \rightarrow \Delta_{\mathcal{X}_s}$ is an isomorphism (since we assumed $p \notin \Sigma$), and the right vertical homomorphism is the natural projection $G_k \twoheadrightarrow G_F$, as follows easily from the specialisation theory for fundamental groups of Grothendieck (cf. [SGA1]).

In fact, In the above commutative diagram (6.2), the right square is cartesian, as follows easily from the slimness of Δ_X , and the well-known criterion for good reduction of curves (cf. [Saïdi], Lemma 4.2.2).

6.2. Let k_X , and k_Y , be two p -adic local fields, i.e. both k_X , and k_Y , are finite extensions of \mathbb{Q}_p .

Let X (resp. Y) be a proper, smooth, geometrically connected, and hyperbolic curve over k_X (resp. k_Y).

Let $\Sigma \subseteq \mathfrak{Primes}$ be a non-empty set of prime integers, and Π_X (resp. Π_Y) the geometrically pro- Σ arithmetic fundamental group of X (resp. Y), which sits in the exact sequence $1 \rightarrow \Delta_X \rightarrow \Pi_X \rightarrow G_{k_X} \rightarrow 1$ (resp. $1 \rightarrow \Delta_Y \rightarrow \Pi_Y \rightarrow G_{k_Y} \rightarrow 1$).

Let

$$\sigma : \Pi_X \xrightarrow{\sim} \Pi_Y$$

be an isomorphism between profinite groups.

Lemma 6.3. *The isomorphism σ fits into a commutative diagram:*

$$(6.3) \quad \begin{array}{ccc} \Delta_X & \longrightarrow & \Delta_Y \\ \downarrow & & \downarrow \\ \Pi_X & \xrightarrow{\sigma} & \Pi_Y \\ \downarrow & & \downarrow \\ G_{k_X} & \longrightarrow & G_{k_Y} \end{array}$$

where the horizontal maps are isomorphisms, which are naturally induced by σ .

In particular, the isomorphism σ induces naturally a bijection

$$\sigma^{\text{sec}} : \overline{\text{Sec}}_{\Pi_X} \xrightarrow{\sim} \overline{\text{Sec}}_{\Pi_Y}.$$

Moreover, the natural isomorphism $G_{k_X} \xrightarrow{\sim} G_{k_Y}$ which is induced by σ preserves the inertia subgroups, i.e. maps the inertia subgroup of G_{k_X} isomorphically to the inertia subgroup of G_{k_Y} .

Proof. (cf. [Mochizuki4], Proposition 1.2.1, and Lemma 1.3.8). \square

Definition 6.4. We say that the isomorphism σ is point-theoretic, if the image of $X(k)$ in $\overline{\text{Sec}}_{\Pi_Y}$, via the map $\sigma^{\text{sec}} \circ \varphi_{X,\Sigma} : X(k) \rightarrow \overline{\text{Sec}}_{\Pi_Y}$, coincides with $\varphi_{Y,\Sigma}(Y(k_Y))$. In other words σ is point-theoretic if it induces naturally a bijection $\varphi_{X,\Sigma}(X(k_X)) \xrightarrow{\sim} \varphi_{Y,\Sigma}(Y(k_Y))$.

It is natural, in the framework of the p -adic GASC, to consider the following question.

Question 6.5 (A Weak Form of the Grothendieck Anabelian Section Conjecture over p -adic Local Fields). Let k_X , and k_Y , be two p -adic local fields. Let X (resp. Y) be a proper, smooth, geometrically connected, and hyperbolic curve over k_X (resp. k_Y).

Assume that $\Sigma = \mathfrak{Primes}$. Let Π_X (resp. Π_Y) be the geometrically pro- Σ fundamental group of X (resp. Y), which sits in the exact sequence $1 \rightarrow \Delta_X \rightarrow \Pi_X \rightarrow G_{k_X} \rightarrow 1$ (resp. $1 \rightarrow \Delta_Y \rightarrow \Pi_Y \rightarrow G_{k_Y} \rightarrow 1$). Let

$$\sigma : \Pi_X \xrightarrow{\sim} \Pi_Y$$

be an isomorphism between profinite groups. Is σ point-theoretic, in the sense of Definition 6.4?

Note that the validity of the p -adic GASC implies a positive answer to Question 6.5.

Although the pro- Σ version of the Grothendieck anabelian section conjecture may not hold over p -adic local fields, in the case where $p \notin \Sigma$ (cf. [Saïdi], Proposition 4.2.1), one may ask whether the following weak form of the pro- Σ Grothendieck anabelian section conjecture still holds, if $p \notin \Sigma$.

Question 6.6. Let k_X , and k_Y , be two p -adic local fields. Let X (resp. Y) be a proper, smooth, geometrically connected, and hyperbolic curve over k_X (resp. k_Y).

Let $\Sigma \subset \mathfrak{Primes}$ be a non-empty set of prime integers, with $p \notin \Sigma$. Let Π_X (resp. Π_Y) be the geometrically pro- Σ fundamental group of X (resp. Y). Let

$$\sigma : \Pi_X \xrightarrow{\sim} \Pi_Y$$

be an isomorphism between profinite groups. Is σ point-theoretic?

Remarks 6.7. A positive answer to Question 6.6 will imply an absolute version of the Grothendieck anabelian conjecture for smooth, proper, and hyperbolic curves over p -adic local fields (cf. [Mochizuki5], Corollary 2.9).

In connection with Question 6.6, we can prove the following.

Proposition 6.8. *Let k_X , and k_Y , be two p -adic local fields. Let X (resp. Y) be a proper, smooth, geometrically connected, and hyperbolic curve over k_X (resp. k_Y).*

Let $\Sigma \subset \mathfrak{Primes}$ be a non-empty set of prime integers, with $p \notin \Sigma$. Let Π_X (resp. Π_Y) be the geometrically pro- Σ fundamental group of X (resp. Y). Let

$$\sigma : \Pi_X \xrightarrow{\sim} \Pi_Y$$

be an isomorphism between profinite groups.

Assume that X (or Y) has good reduction over k_X (or k_Y), i.e. X (or Y) extends to a proper, and smooth, relative curve over the valuation ring O_{k_X} of k_X (or, over the valuation ring O_{k_Y} of k_Y). Then σ is point-theoretic.

Proof. First, X has good reduction over k_X if and only if Y has good reduction over k_Y , as follows easily from the well-known criterion for good reduction, and the last assertion in Lemma 6.3.

Assume that X (resp. Y) extends to a proper and smooth relative curve \mathcal{X} over the valuation ring O_{k_X} of k_X (resp. \mathcal{Y} over the valuation ring O_{k_Y} of k_Y). Let \mathcal{X}_s (resp. \mathcal{Y}_s) be the special fiber of \mathcal{X} (resp. \mathcal{Y}).

Recall the commutative diagrams (6.2):

$$\begin{array}{ccccccc} 1 & \longrightarrow & \Delta_X & \longrightarrow & \Pi_X & \longrightarrow & G_{k_X} \longrightarrow 1 \\ & & \downarrow & & \text{Sp}_X \downarrow & & \downarrow \\ 1 & \longrightarrow & \Delta_{\mathcal{X}_s} & \longrightarrow & \Pi_{\mathcal{X}_s} & \longrightarrow & G_{F_X} \longrightarrow 1 \end{array}$$

and

$$\begin{array}{ccccccc} 1 & \longrightarrow & \Delta_Y & \longrightarrow & \Pi_Y & \longrightarrow & G_{k_Y} \longrightarrow 1 \\ & & \downarrow & & \text{Sp}_Y \downarrow & & \downarrow \\ 1 & \longrightarrow & \Delta_{\mathcal{Y}_s} & \longrightarrow & \Pi_{\mathcal{Y}_s} & \longrightarrow & G_{F_Y} \longrightarrow 1 \end{array}$$

where F_X (resp. F_Y) is the residue field of k_X (resp. k_Y).

Let $s = s_x : G_{k_X} \rightarrow \Pi_X$ be a group-theoretic section of the natural projection $\Pi_X \twoheadrightarrow G_{k_X}$, which is point-theoretic, i.e. arises from a rational point $x \in X(k_X)$.

Then s induces naturally a group-theoretic section $s' : G_{k_Y} \rightarrow \Pi_Y$ of the natural projection $\Pi_Y \twoheadrightarrow G_{k_Y}$, such that the following diagram is commutative:

$$\begin{array}{ccc} G_{k_X} & \longrightarrow & G_{k_Y} \\ s \downarrow & & s' \downarrow \\ \Pi_X & \xrightarrow{\sigma} & \Pi_Y \end{array}$$

where the upper arrow is the natural isomorphism which is induced by σ . We will show that s' is point-theoretic.

The natural map $\mathrm{Sp}_X \circ s_x : G_{k_X} \rightarrow \Pi_{\mathcal{X}_s}$ factorises as $G_{k_X} \twoheadrightarrow G_{F_X} \xrightarrow{\bar{s}} \Pi_{\mathcal{X}_s}$, where $\bar{s} : G_{F_X} \rightarrow \Pi_{\mathcal{X}_s}$ is a group-theoretic section of the natural projection $\Pi_{\mathcal{X}_s} \twoheadrightarrow G_{F_X}$.

The section \bar{s} is point-theoretic since s is, and corresponds to a decomposition subgroup associated to the point $\bar{x} \in \mathcal{X}_s(F_X)$, which is the specialisation of the rational point $x \in X(k_X)$.

Let

$$\dots \subseteq \Delta_X[i+1] \subseteq \Delta_X[i] \subseteq \dots \subseteq \Delta_X[1] \stackrel{\mathrm{def}}{=} \Delta_X$$

be a family of open characteristic subgroups of Δ_X , with $\bigcap_{i \geq 1} \Delta_X[i] = \{1\}$. We also denote by $\{\Delta_X[i]\}_{i \geq 1}$ the corresponding family of open subgroups of $\Delta_{\mathcal{X}_s}$, via the specialisation isomorphism $\mathrm{Sp}_X : \Delta_X \xrightarrow{\sim} \Delta_{\mathcal{X}_s}$.

For a positive integer i , let $\Pi_X[s, i] \stackrel{\mathrm{def}}{=} \Delta_X[i].s(G_{k_X})$, and $\Pi_{\mathcal{X}_s}[\bar{s}, i] \stackrel{\mathrm{def}}{=} \Delta_X[i].\bar{s}(G_{F_X})$. The system of open subgroups $\{\Pi_X[s, i]\}_{i \geq 1}$ (resp. $\{\Pi_{\mathcal{X}_s}[\bar{s}, i]\}_{i \geq 1}$) correspond to a tower of finite étale covers

$$\dots \rightarrow X_{i+1} \rightarrow X_i \rightarrow \dots \rightarrow X_1 \stackrel{\mathrm{def}}{=} X,$$

(resp.

$$\dots \rightarrow \mathcal{X}_{s,i+1} \rightarrow \mathcal{X}_{s,i} \rightarrow \dots \rightarrow \mathcal{X}_{1,s} \stackrel{\mathrm{def}}{=} \mathcal{X}_s),$$

where $\{X_i\}_{i \geq 1}$ (resp. $\{\mathcal{X}_{s,i}\}_{i \geq 1}$) form a system of neighbourhoods of the section s (resp. \bar{s}).

Let $\Pi_Y[i] \stackrel{\mathrm{def}}{=} \sigma(\Pi_X[s, i])$, and $\Pi_{\mathcal{Y}_s}[i] \stackrel{\mathrm{def}}{=} \sigma(\Pi_{\mathcal{X}_s}[\bar{s}, i])$. The natural map $\mathrm{Sp}_Y \circ s' : G_{k_Y} \rightarrow \Pi_{\mathcal{Y}_s}$ factorises as $G_{k_Y} \twoheadrightarrow G_{F_Y} \xrightarrow{\bar{s}'} \Pi_{\mathcal{Y}_s}$, where $\bar{s}' : G_{F_Y} \rightarrow \Pi_{\mathcal{Y}_s}$ is a group-theoretic section of the natural projection $\Pi_{\mathcal{Y}_s} \twoheadrightarrow G_{F_Y}$ (this factorisation is induced by the similar factorisation of $\mathrm{Sp}_X \circ s$).

The $\{\Pi_Y[i]\}_{i \geq 1}$ (resp. $\{\Pi_{\mathcal{Y}_s}[i]\}_{i \geq 1}$) form a system of neighbourhoods of the section s' (resp. \bar{s}'), and correspond to a tower of étale covers

$$\dots \rightarrow Y_{i+1} \rightarrow Y_i \rightarrow \dots \rightarrow Y_1 \stackrel{\mathrm{def}}{=} Y$$

(resp.

$$\dots \rightarrow \mathcal{Y}_{s,i+1} \rightarrow \mathcal{Y}_{s,i} \rightarrow \dots \rightarrow \mathcal{Y}_{s,1} \stackrel{\mathrm{def}}{=} \mathcal{Y}_s).$$

The section \bar{s}' is naturally induced by the section \bar{s} via the natural isomorphism $\Pi_{\mathcal{X}_k} \xrightarrow{\sim} \Pi_{\mathcal{Y}_k}$. In particular, the section \bar{s}' is point-theoretic, since \bar{s} is point-theoretic, as follows from the arguments of Tamagawa for characterising decomposition groups of rational points over finite fields (see for example [Saïdi-Tamagawa], §1), hence $\mathcal{Y}_{s,i}(F_Y) \neq \emptyset$.

Also, by the very definition of the system of neighborhoods $\{Y_i\}_{i \geq 1}$, each Y_i has good reduction over \mathcal{O}_{k_Y} , and extends to a smooth and proper model over \mathcal{O}_{k_Y} whose special fibre is $\mathcal{Y}_{s,i}$. In particular, $Y_i(k_Y) \neq \emptyset$ by the theorem of liftings of smooth points (cf. [SGA1], exposé III, Corollaire 3.3). This implies that the section s' is point-theoretic by Lemma 2.5. \square

In connection to question 6.6, and under the condition that the curve X (or Y) has potentially good reduction, one may hope to use Proposition 6.8, plus a descent argument, to give a positive answer to question 6.6 in this case. The author has no idea, at the moment of writing this paper, on how to perform such a descent argument.

References.

- [Esnault-Wittenberg] Esnault, H., Wittenberg, O., Remarks on cycle classes of sections of the fundamental group, *Mosc. Math. J.* 9 (2009), no. 3, 451-467.
- [Esnault-Wittenberg1] Esnault, H., Wittenberg, O., On abelian birational sections. *Journal of the American Mathematical society*, Volume 23, Number 3, July 2010, Pages 713-724.
- [Grothendieck] Grothendieck, A., Brief an G. Faltings, (German), with an english translation on pp. 285-293. *London Math. Soc. Lecture Note Ser.*, 242, Geometric Galois actions, 1, 49-58, Cambridge Univ. Press, Cambridge, 1997.
- [Hoshi] Hoshi, Y., Existence of nongeometric pro- p Galois sections of hyperbolic curves. To appear in *Publications of RIMS*.
- [Koenigsmann] Koenigsmann, J., On the section conjecture in anabelian geometry. *J. Reine Angew. Math.* 588 (2005), 221-235.
- [Mochizuki] Mochizuki, S., Absolute anabelian cuspidalizations of proper hyperbolic curves. *J. Math. Kyoto Univ.* 47 (2007), no. 3, 451-539.
- [Mochizuki1] Mochizuki, S., The local pro- p anabelian geometry of curves. *Invent. Math.* 138 (1999), no. 2, 319-423.
- [Mochizuki2] Mochizuki, S., Topics surrounding the anabelian geometry of hyperbolic curves. *Galois groups and fundamental groups*, 119-165, *Math. Sci. Res. Inst. Publ.*, 41, Cambridge Univ. Press, Cambridge, 2003.
- [Mochizuki3] Mochizuki, S., Galois sections in absolute anabelian geometry. *Nagoya Math. J.* 179 (2005), 17-45.
- [Mochizuki4] Mochizuki, S., The absolute anabelian geometry of hyperbolic curves. *Galois theory and modular forms*, 77-122, *Dev. Math.*, 11, Kluwer Acad. Publ., Boston, MA, 2004.
- [Mochizuki5] Mochizuki, S., Topics in absolute anabelian geometry II: decomposition groups and endomorphisms. Preprint. Available in the home web page of Shinichi Mochizuki.
- [Pop] Pop, F., On the birational p -adic section Conjecture, *Compos. Math.* 146 (2010), no. 3, 621-637.
- [Saïdi] Saïdi, M., Good sections of arithmetic fundamental groups. Manuscript.
- [Saïdi-Tamagawa] Saïdi, M., Tamagawa, A., A prime-to- p version of the Grothendieck anabelian conjecture for hyperbolic curves over finite fields of characteristic $p > 0$. *Publ. Res. Inst. Math. Sci.* 45 (2009), no. 1, 135-186.
- [SGA1] Grothendieck, A., Revêtements étales et groupe fondamental. *Séminaire de géométrie algébrique du Bois Marie 1960-61*.

[Tamagawa] Tamagawa, A., The Grothendieck conjecture for affine curves. *Compositio Math.* 109. (1997), no. 2, 135–194.

Mohamed Saïdi
College of Engineering, Mathematics, and Physical Sciences
University of Exeter
Harrison Building
North Park Road
EXETER EX4 4QF
United Kingdom
M.Saïdi@exeter.ac.uk